

Dopamine Reward Learning Models and Decision Making Under Uncertainty

Rama Khan^{1,2}, Pakiza Ikram³, Nimra Hameed⁴

¹Department of Nutrition and Dietetics, University of Faisalabad, Faisalabad, Pakistan; ²Department of Nutrition and Dietetics, Istanbul Gelisim University, Istanbul, Türkiye (Erasmus Exchange Student); ³Department of Computer Science, The Superior College Lahore, Pakistan; ⁴Department of Computer Science, Virtual University of Pakistan

ABSTRACT

Dopamine reward learning models make a substantial contribution to the description of how individuals and animals make decisions in ambiguous situations. These models are primarily based on reinforcement learning. This theory further posits that dopamine neurons encode reward prediction errors, defined as the difference between expected and actual outcomes. Such signals enable the brain to update its value representations and to guide future decision-making in dynamic, unpredictable environments. Dopamine-mediated learning helps maintain a healthy balance between exploration and exploitation in uncertain situations by modulating the sensitivity of risk, motivation, and behavioural flexibility. It has been shown that dopaminergic signals are adapted to probabilistic incentives, delayed information, and ambiguous information using computational frameworks. These models are the temporal-difference learning and the Bayesian decision models. Maladaptive decision-making has been observed to be linked to disruptions in dopamine signalling, including schizophrenia, Parkinson's, and addiction, which are the results of empirical research in the fields of neuroscience, psychology, and neuroeconomics. Dopamine reward learning provides an excellent account of how to process uncertainty and make optimal or suboptimal decisions. It is achieved through combining computer models with brain mechanisms. This fact not only advances cognitive neuroscience but also provides information for artificial intelligence systems designed to operate in uncertain and complex environments.

ARTICLE HISTORY

Received September 10, 2025

Accepted November 18, 2025

KEYWORDS

Dopamine Reward (DR),
Learning Models (LM),
Decision Making (DM),
Uncertainty (UC)

1. Introduction

Human beings continually face situations that require learning from experience and making choices in environments where outcomes are uncertain. In daily life, we encounter countless examples of this process: choosing the best route to work, deciding whether to trust new information, selecting financial options, or even forming social judgments. These tasks depend on an internal ability to evaluate rewards, estimate future consequences, and modify actions in response to changing circumstances (Ban, 2024). These abilities are neural mechanisms that understand events, form predictions, and update beliefs. Over time, extensive research has converged on the view that dopamine plays a central role in reward-based learning and adaptive decision-making, particularly under conditions of uncertainty (Ben-Zion & Levy, 2025). Learning from experience is often explained by reinforcement principles. When we obtain desirable outcomes by taking a particular action, the likelihood of repeating that action increases; when the outcome is disappointing, the behaviour is adjusted. This cycle depends on comparing expected outcomes with actual outcomes (Ghosh et al., 2025). The difference between them is known as prediction error, and it serves as a teaching signal that helps to improve future decisions. Dopamine activity aligns closely with this process. Dopamine neurons increase their firing when consequences exceed expectations and reduce firing when expected rewards do not appear (Skowron, 2024). The brain consistently recalibrates its internal expectations and action strategies through this dynamic signalling. Furthermore, computational models of reinforcement learning have provided a powerful framework to understand these processes. The first algorithm in this field is called temporal-difference learning, which adjusts choices based on the difference between anticipated and received rewards at each time step (Paunov et al., 2024). The dopamine activity maps up to this algorithm with extraordinary precision, providing a biological explanation for the mathematical form of reinforcement learning. This correspondence has positioned dopamine as more than a chemical associated with reward, as it functions as a central element of neural

computation systems, facilitating flexible learning (Korbisch & Ahmed, 2025).

Decision-making becomes more complex in environments where outcomes and rewards are not fully predictable. Uncertainty can arise from an inherently variable environment, imperfect access to information, or rapid changes in conditions over time. To work effectively in these environments, the brain must assess the reliability of past experiences and determine how strongly new information and knowledge should influence behaviour (Story et al., 2024). This process involves not only estimating reward value but also weighing probabilities, detecting changes, and adjusting learning rates. Dopamine contributes to these adjustments by shaping the weight assigned to a new outcome (Taira & Sharpe, 2025). In addition, small learning updates help maintain consistent behaviour in a stable environment, whereas larger updates promote rapid adaptation in a rapidly changing environment. The variation of learning rates allows an individual to manage stability with flexibility. For example, when unexpected outcomes recur, dopamine signalling can enhance sensitivity to new information, thereby encouraging exploration of alternative actions. Dopamine in a predictable environment facilitates the formation of established habits by reinforcing existing expectations (Attaallah et al., 2024). This adaptive response links dopamine to a broader problem of action under uncertainty. Dopamine affects various parts of the brain, which contribute to our decision-making. The ventral striatum is primarily an indication of anticipatory rewards and inappropriate alterations of these rewards when they are not attained. The prefrontal cortex helps us plan, evaluate options, and reason about long-term consequences (Feng et al., 2024). Ambiguous decision-making is a major challenge that human beings and animals face in their day-to-day lives. Humans are frequently unable to make fully informed decisions about what to invest in, which professions to pursue, and how to respond to environmental threats. Across neuroscience, psychology, and cognitive science, one of the most significant questions concerns how the brain acquires knowledge through experience and how it alters its behaviour in uncertain circumstances. Among the various neurobiological processes involved, dopamine-based reward-learning models have become among the most prominent frameworks for explaining adaptive decision-making. Dopamine is a neuromodulator that is extremely crucial in motivation, learning, and reward processing. Initial research linked dopamine solely to pleasure and reward, but current research indicates a more complex role. The dopaminergic neurons, especially the ones found in the midbrain, like the ventral tegmental area (VTA) and the substantia nigra, encode the reward prediction errors- the difference between the provided and obtained results. This cue is an instructional mechanism that enables the brain to adjust expectations and future action. This kind of learning is critical in an uncertain environment in which outcomes are probabilistic, delayed, or vague. Reinforcement learning (RL) models are based on the idea of predicting a reward signal and explaining how agents learn optimal behaviour through trial and error. Computational view: Temporal-difference learning provides reinforcement-learning paradigms that offer a computationally grounded understanding of the role of dopamine signals in learning and decision-making. These models suggest that dopamine activity increases when outcomes exceed predictions, whereas it decreases when outcomes fall below predictions. Over time, these signals develop representations of values associated with actions or stimuli, thereby enabling individuals to make informed decisions. Most realistic decision situations are characterized by uncertainty. In contrast to deterministic environments, uncertain environments are characterized by imperfect information, variable reward systems and variable situations. The process of decision-making under uncertainty requires humans to make probability estimates, accept risk, and adapt to unforeseen consequences. The anterior cingulate cortex is involved in confronting and struggling with environmental changes. The integration of these areas enables the combination of reward signals, uncertainty assessments, and goals to inform action (Cecchini et al., 2024). Learning, motivation, and evaluation of all these parts are coordinated by dopamine. This model helps us to explain several human actions. When the environment becomes uncertain, individuals become more inquisitive, seeking alternative solutions until a consistent pattern emerges. As circumstances stabilise, behaviour becomes more predictable and routine. These dynamics are based on how dopamine interacts to generate prediction-error signals and uncertainty-based adjustments (Nussenbaum & Hartley, 2024). Dopamine system failure can cause serious problems with learning and decision-making. Parkinson's disease creates low dopamine levels, which decrease good learning results and increase sensitivity to unfavourable consequences. This imbalance reduces motivation and flexibility.

During addiction, reward systems become hyperresponsive to signals linked with dangerous drugs, rewarding poor habits over good ones. Disorders such as schizophrenia and obsessive-compulsive disorder both exhibit dopamine-mediated altered beliefs, risk evaluation, and adaptability (Dillon et al., 2024). These examples illustrate how dopamine-based learning frameworks influence both healthy and unhealthy choices. Recent advances in neuroimaging, computational modelling, and electrophysiology have made it clear that dopamine plays an important role in decision-making under uncertainty. Brain imaging studies typically show dopamine-related activity during tasks that require predicting rewards, estimating likelihoods, adjusting behaviour, and responding to unexpected events (Mah et al., 2024). The study of reward learning and uncertainty also involves the development of internal representations of the world in the brain. These models enable people to make predictions, plan about what is likely to happen, and understand emerging data. Dopamine helps update the models when predictions are missed or when conditions change. Individuals improve their ability to recognise

patterns, adverse consequences, and motivation with time (Wheeler et al., 2024). In uncertain environments, accurate models are even more important, since even small prediction errors can lead to undesirable outcomes. Dopamine, therefore, serves as a key signal that keeps the models consistent with reality. Overall, the dopamine-reward learning model provides a useful framework for understanding the mechanisms underlying behavioural adaptation in unpredictable and changing environments. The complex relationship between dopamine signalling and reinforcement learning explains why people recognise imbalances, reprioritise, and adjust their behaviour.

2. Literature Review

Uncertain decision-making has become a major subject in neuroscience and computational psychology. Most research focuses on the role of dopamine in reward-directed behaviour. The processes of recording prediction errors, estimating value, and learning-rate adaptability in dynamic situations have been investigated over time in relation to dopamine circuits (Liebana et al., 2024). Studies have shown that dopamine facilitates the acquisition of reward information and the refinement of behaviour in response to unexpected outcomes. Researchers reported that computer models, human imaging experiments, animal research, and clinical investigations collectively illuminate the function of dopamine in uncertainty-motivated behaviour (Blanco-Pozo et al., 2024). It was examined that one of the primary areas of study was the relationship between dopamine and reward error prediction. Studies have shown that dopamine-producing neurons produce greater output when outcomes exceed expectations and less when expected rewards are absent. Such a firing pattern supports temporal-difference learning theories, which posit that behaviour is modified by the difference between actual and predicted outcomes (Gershman & Lak, 2025). Activities with probabilistic incentives show that the signal of prediction error was related to uncertainty. As the probability of a reward was varied, dopamine activity became more sensitive to aversive surprises, underscoring the need for greater flexibility in adaptation. Researchers highlighted that another key question concerns how dopamine regulates learning rates in both variable and stable contexts (Kahnt & Schoenbaum, 2025). Lower levels of learning have been utilised in stable conditions to maintain a consistent set of expectations. In contrast, higher learning rates have been used in unstable environments to adapt quickly. Tests of reinforcement learning under shifting reward conditions show that dopamine levels signal environmental uncertainty. According to many researchers, high levels of dopaminergic activity enhance learning and enable rapid updating of results in response to environmental instability (Metha et al., 2025). As a result, dopamine governs the brain's susceptibility to novel information and also serves as a reward signal. Researchers explored how the brain generates expectations about the desirability of acts or outcomes, rather than focusing just on prediction errors. The ventral striatum and prefrontal cortex receive dopaminergic projections that encode predicted value, enabling individuals to choose the activity with the highest reward value.

Neuroimaging studies of the human brain during dangerous decision-making indicate increased activity in these regions when individuals consider questionable options (Gershman et al., 2024). These findings suggest that the dopamine circuits that determine subjective value consider both the magnitude and the possibility of reward. The exploration-exploitation dilemma asks whether one should continue a known, reward-predictable behaviour (exploitation) or pursue unexplored possibilities. Dopamine helps resolve this by influencing novelty-seeking behaviour and various habits, thereby shifting decisions toward exploratory ones (Schultz, 2024). Studies show that increased dopaminergic tone correlates with increased exploratory behaviour under unpredictable reward schedules. It was indicated that dopamine facilitates the adaptive processes necessary to cope with novel environments by promoting behavioural variability. The other key strand focuses on the function of dopamine in risk preferences (Kurtenbach et al., 2024). Dopamine reward learning models provide an attractive account of how the brain overcomes these obstacles. Dopaminergic systems help decision-makers remain flexible and responsive to environmental changes by dynamically adjusting value expectations in response to prediction errors, according to prediction errors. The trade-off between exploration and exploitation is a major problem in decision-making under uncertainty. Exploration involves considering less familiar or new options to seek information, whereas exploitation involves pursuing available, familiar options to reap rewards. Dopamine is a key factor in regulating this balance. The high level of dopamine activity has been associated with exploration and risk-taking, and the low levels of dopamine have been associated with conservative and exploitative behaviour. Computational models suggest that the rates of learning and uncertainty sensitivity are regulated by dopamine, such that one can adjust one's approach to environmental volatility. Alongside reinforcement learning models, Bayesian decision-making models have been combined with dopamine-based learning theories to represent uncertainty more accurately. Bayesian models emphasize updating beliefs using prior information and newly received data, and provide a probabilistic perspective on learning and decision-making. It has been suggested that dopamine signals may be involved in reward prediction errors, as well as in confidence, uncertainty, and belief revision. This combination has enhanced our understanding of how the brain processes uncertainty and adapts behavior in complex and confusing environments. Much of the empirical evidence of dopamine reward learning models is based on a diverse variety of experimental paradigms. Neurophysiological studies in animals indicate that

dopamine neurons respond to unexpected rewards and to reward anticipation. Neuroimaging research on the human brain also shows that these dopaminergic regions are activated during gratifying learning and risky decision-making. According to behavioural tests, human beings adjust their choices in response to feedback in a manner that closely aligns with predictions from reinforcement learning models. Researchers have demonstrated that risky decision-making involves evaluating potential outcomes in terms of probabilities. Dopamine regulates how people consider the magnitudes of rewards against the chances. According to drug studies, increasing dopamine can increase sensitivity to potential rewards, sometimes leading to a risk-seeking reaction, whereas decreasing dopamine switches preferences to less harmful choices (Saxena, 2024). Therefore, changes in dopamine signalling significantly affect decision-making in ambiguous situations. Recent research investigated the role of dopamine in updating beliefs under severe uncertainty, including volatility and ambiguity. Ambiguity concerns uncertainty about probabilities, whereas volatility refers to unpredictability in a changing environment. Dopamine helps detect changes in environmental structure and modulate learning strategies (Bruckner & Nassar, 2024).

According to computational theories, dopamine signals underlie the unpredictability of events. This investigation evaluates the extent to which new evidence influences individuals' opinions. As a result, dopamine plays a significant role in metacognition and in the desire to improve adaptive learning (Augustat et al., 2024). Clinical studies have shown dopamine's functions in a variety of diseases, including addiction, schizophrenia, Parkinson's disease, and obsessive-compulsive disorder, among others. Researchers highlighted that reward signals generate excessive dopaminergic reactivity in people with an addiction, which promotes harmful behaviours. Parkinson's patients have low amounts of dopamine, which inhibits consecutive learning based on good results while encouraging learning based on negative consequences. Dopaminergic processing problems lead to inflexible or maladaptive decision-making, demonstrating that clinical information complements computational models of behaviour. According to many researchers, activating dopamine neurons during reward-related activities can artificially enhance learning, whereas preventing their activation disrupts incentive-driven behaviour (Khan, 2025). Animal studies reveal that dopamine interacts with other neuromodulators, such as serotonin and norepinephrine, to modulate the processing of uncertainty. As a result, it was observed that dopamine's contribution to decision-making arises from its interaction with broader brain circuits rather than from autonomous action. Human neuroimaging has been essential for revealing the involvement of dopamine-rich regions in the performance of uncertain tasks. Functional MRI consistently shows that the striatum, prefrontal cortex, and midbrain are activated during probabilistic judgment (Dias Maile et al., 2025). These activations have been associated with computational signals, such as prediction inaccuracy, anticipated value, and risk estimate. Researchers examined diffusion imaging, which demonstrates that the structural connectivity of dopaminergic pathways and cortical regions predicts individual variation in uncertain-reward learning (Wang et al., 2024). These tendencies suggest that anatomical and functional aspects of dopamine circuits influence how people understand uncertainty and, in turn, how they behave. Finally, scholars have shown that cognitive characteristics related to dopamine (e.g., reward sensitivity or learning-rate asymmetry) may differ among individuals, and these differences are components of individual decision-making. Behavioural preferences may be classified using computational indicators of dopaminergic activity, which could enable future diagnostic and therapeutic breakthroughs (Chou et al., 2025). Critically, shifts in dopamine signaling provide a novel perspective on dopamine's involvement in decision-making under uncertainty. The clinical conditions are linked with impaired dopamine activity and decreased decision-making capacity, such as Parkinson's disease, addiction, attention-deficit/hyperactivity disorder (ADHD), and schizophrenia. As an illustration, maladaptive learning on positive consequences in response to activities like dopamine loss in Parkinson's disease, and positive activity of dopamine in addiction, are linked to maladaptive reward-seeking behavioural patterns despite the negative consequences. These results highlight the delicate equilibrium required for effective dopaminergic function and adaptive decision-making. Outside of neuroscience and psychology, dopamine reward-learning models have had a significant influence on artificial intelligence and machine learning. Biological dopamine-based reinforcement learning algorithms have now become common in robotics, game-playing agents, and adaptive control systems. Such biologically inspired models enable artificial agents to learn from feedback and make decisions under ambiguous conditions, thereby mimicking basic properties of human and animal learning. Consequently, the dopamine-based theory has been used to connect biological and artificial intelligence. However, despite such enormous progress, several obstacles remain to the proper interpretation of the role of dopamine in decision-making under uncertainty. The dopaminergic system interacts with other brain regions, such as the prefrontal cortex, striatum, and limbic system, and its activity is highly context dependent.

Furthermore, dopamine signals can reflect more than fundamental reward prediction errors, such as motivational salience or uncertainty itself. Current research continues to refine earlier models and elucidate the role of dopamine in interacting with other neuromodulator systems in the process of making complex decisions. In short, dopamine reward learning models provide a comprehensive account of how decisions are formed in the face of uncertainty. These models explain how prediction errors drive learning, how uncertainty affects behavior, and why decision-making is prone to

maladaptive behaviour when dopamine signaling is disrupted. As research continues, dopamine-based models will remain central to advancing our understanding of cognition, behaviour, and intelligent systems in uncertain situations (Figure 1).




Figure 1: Time-Series Comparison of Neurotransmitter Levels

3. Methodology

The analysis presented here employs a hybrid theoretical-computational approach to assess the importance of dopamine-based reward-learning models in uncertain decision-making. The method combines computer simulation, analysis of behavioural tasks, and secondary analysis of neuroscientific evidence to provide comprehensive information on the mechanisms of learning involving dopaminergic systems. A reinforcement learning (RL) theory of computational modelling is initially used. Temporal-Difference (TD) learning models simulate errors in reward prediction caused by dopamine. Systematic variations in model parameters, such as the learning rate, discount factor, and exploration parameter, are used to investigate the effects of dopamine signalling on learning and choice behaviour in an uncertain reward environment.

3.1 Dopamine Reward Learning Models

Deterministic and probabilistic reward systems representing varying degrees of environmental uncertainty are simulated. Second, standardised behavioural paradigms commonly used in neuroscience studies are employed to investigate decision-making. Computational representation, Probabilistic reward learning, the problem of multi-armed bandits, and risk-based choice problems are discussed. These tasks enable one to test exploration-exploitation trade-offs, sensitivity to rewards and adaptation to new reward conditions. Predictions from the RL models are compared with behavioural results to evaluate their explanatory capability. Third, neurobiological evidence is incorporated, based on a systematic review of empirical studies conducted by peers. Neuroimaging and neurophysiological studies in humans and animal models are examined to support computational predictions about dopamine reward errors. Specific emphasis is given to studies that examine midbrain dopamine activity, striatal value encoding, and prefrontal control mechanisms in the context of ambiguity. The models' performance is assessed using measures of prediction accuracy, likelihood estimation, and goodness of fit. To assess the models' robustness to variation in uncertainty and reward volatility, sensitivity analyses are performed. This triangulated methodological design will ensure that theoretical, computational, and empirical perspectives are consistently considered, thereby enhancing the validity and reliability of the results (Figure 2).

Reversal learning




Figure 2: Dopamine Activity Measurement in a Reversal Learning Task

4. Equational Models

Dopamine reward learning equational models are derived primarily from reinforcement learning theory and provide a formal mathematical account of how agents learn under conditions of uncertainty and reward inputs. According to these models, dopamine neurons encode errors in reward prediction, which are used to reshape value estimates in relation to states or behaviors.

4.1 Reward Prediction Error

The fundamental model in dopamine-based learning models is the reward prediction error (RPE), which is expressed as:

$$\delta_t = r(s(t+1)) - V(s_t)$$

where

δ_t represents the reward prediction error at time t ,

r_t is the reward received,

γ is the discount factor ($0 \leq \gamma \leq 1$),

$V(s_t)$ and $V(s_{t+1})$ denote the value of the current and subsequent states, respectively.

A positive δ_t indicates that outcomes are better than expected, leading to increased dopamine activity, while a negative δ_t reflects worse-than-expected outcomes and reduced dopamine signalling.

4.2 Value Update Rule

Value estimates are updated using a learning rule based on the prediction error:

$$V(s_t) \leftarrow V(s_t) + \alpha \delta_t$$

Where α is the learning rate, ($0 < \alpha \leq 1$). Higher learning rates correspond to faster updating, often associated with heightened dopaminergic sensitivity, while lower rates reflect more conservative learning under uncertainty.




Figure 3: Q-Learning vs. Deep Q-Learning for Action-Value Estimation

4.3 Action-Value (Q-Learning) Model

When decisions involve choosing among multiple actions, action-value learning is used (Figure 3):

$$Q(s_t, a_t) \leftarrow Q(s_t, a_t) + \alpha \left[r_t + \gamma \max_a Q(s_{t+1}, a) - Q(s_t, a_t) \right]$$

This formulation allows the agent to learn which actions yield higher expected rewards in uncertain environments.

4.4 Action Selection Under Uncertainty

Decision-making is modelled using a SoftMax action selection rule:

$$P(a_t | s_t) = \frac{e^{\beta Q(s_t, a_t)}}{\sum_a e^{\beta Q(s_t, a)}}$$

Where β is the inverse temperature parameter that controls the exploration–exploitation balance. Lower β values increase exploratory behaviour, whereas higher values favour the exploitation of known rewarding actions.

4.5 Modeling Uncertainty and Risk

Uncertainty is incorporated by allowing learning rates or decision parameters to vary with environmental volatility:



$$\alpha_t = f(\sigma_t)$$

Where σ_t This represents estimated uncertainty; this dynamic adjustment reflects empirical findings that dopamine signalling adapts to changes in reward predictability.

4.6 Bayesian Integration

In Bayesian reinforcement learning models, belief updating follows:

$$P(\theta | D) \propto P(D | \theta)P(\theta)$$

Where θ represents latent reward parameters and D observed data. Dopamine signals are hypothesized to encode deviations from expected beliefs, integrating probabilistic inference with reward learning. These are equational models that provide a formal description of the interplay between dopamine neurophysiology and measurable decision-making behavior. They provide the mathematical framework of the study of decision-making in uncertainty, which is rigorous by encapsulating prediction errors, value updates, action selection, and adaptation to uncertainty (Figure 4).




Figure 4: Reward Prediction Error Mechanism

5. Discussion

The current argument highlights the importance of dopamine-based reward-learning models in the study of decision-making under uncertain conditions. These models combine neurobiological information with computational models to provide a unified account of learning and adaptive behaviour in settings with limited knowledge, probabilistic outcomes, and dynamic reward schemes. The results presented in this article support the essential role of dopamine-mediated errors in reward prediction, which affects not only short-term decision-making but also long-term behavioural plans. Among the key findings of dopamine reward learning models is that they can explain how individuals adjust their expectations in response to unanticipated outcomes. Reward prediction errors provide a biologically plausible mechanism by which the brain regularly modifies value estimates. This is especially effective in unpredictable situations, and results may not be as expected. The congruence between computational predictions and empirical brain data also enhances the validity of such models and contributes to their applicability across species and contexts. The paper also highlights the importance of dopamine in the regulation of the exploration trade-off. In situations of uncertainty, decision-makers must trade off the urge to seek new options against the desire to harvest known incentives. Dopamine concentrations modify this balance by changing the speed of learning and action-choice strategies. High activity of dopaminergic is associated with increased exploratory behaviour and risk-taking, and low activity is related to cautious and exploitative assessment. Such dynamic adaptation enables people to be flexible in challenging situations, but can equally result in poor decisions in cases where dopamine communication is disorganised (Chou et al., 2025).

Another critical factor is the influence of uncertainty on dopamine response elicitation. Conventional theories view dopamine primarily as a signal of prediction error. However, the data are rapidly accumulating, indicating that dopamine can also serve as an expression of doubt, confidence, or motivational salience. Bayesian reinforcement learning models offer an excellent means of representation of these add-on attributes. Such models provide a more comprehensive account of the brain as an indicator of uncertainty and of how it adjusts behaviour by incorporating belief updating and probabilistic inference. Dopamine reward learning models are also more useful in clinical settings. The disorders like Parkinson's disease, addiction, and schizophrenia accentuate the fact that when abnormalities occur in the process of transmission of dopamine, it hinders the learning process and the process of making decisions under ambiguity. An example is that dopamine loss would reduce sensitivity to reinforcing stimuli. In contrast, excess dopamine signalling would escalate the tendency to seek rewards regardless of their harmfulness. This evidence highlights the sensitivity required for efficient

dopaminergic function and the importance of models that account for both hypo- and hyper-dopaminergic conditions. The dopamine reward-learning models have several disadvantages despite their virtues. Most computational models make simplified assumptions about neural processes and contextual structure and are not necessarily as complex as real-world decision-making. Additionally, dopamine does not act alone; it interacts with other neuromodulator systems, including serotonin and norepinephrine, and with higher-order cognitive functions in the prefrontal cortex. The following generation models should therefore be more integrative in their approach to these relationships. In sum, dopamine reward learning models are an effective and flexible framework for understanding decision-making under uncertainty. Although there are still issues to address in enhancing these models and expanding their explanatory scope, their ability to integrate neurobiological processes, behaviour, and computational theory renders them essential to current research in neuroscience, psychology, and artificial intelligence.

6. Conclusion

This paper concludes that dopamine reward-learning models provide an integrated and detailed framework for explaining decision-making under uncertainty. These models advance understanding of how agents adjust their behaviour in situations characterised by partial information, probabilistic rewards, and dynamics by integrating neurobiology with computational learning theories. The key element of this method is the concept of reward prediction error, which allows the value expectations to be continually revised and facilitates adaptive learning in the long term. The facts presented in this article indicate that dopamine signalling is critical to the direction of the learning process and decision-making behaviour. Dopaminergic neurons encode the difference between anticipated and actual outcomes, enabling individuals to adjust decision-making strategies in response to feedback. It is an effective process, particularly in uncertain conditions in which expectations are often violated. These dynamics can be successfully captured using reinforcement learning models grounded in dopamine dynamics, thereby providing mathematically accurate and biologically plausible explanations of behaviour. One of the strengths of the dopamine-based models is that they explain the trade-off between exploration and exploitation. Ambiguous decision-making requires a balance between seeking new information and exploiting proven, lucrative opportunities. Dopamine regulates this balance by influencing learning rates, reward sensitivity, and risk-taking decisions. With this modulation, human beings remain adaptive and adjust to environmental turbulence. Nevertheless, the same process results in maladaptive decision-making in case dopamine transmission is dysregulated, which occurs in most neurological and psychiatric disorders.

The combination of Bayesian methods of the reinforcement learning model significantly enhances the explanatory capacity of the dopamine reward learning theories. These hybrid model additions of belief updating and probabilistic inference further characterise uncertainty and cognitive flexibility. This has been accompanied by a growing appreciation that dopamine signalling can encode not only reward prediction errors but also aspects of uncertainty, confidence, and motivational salience. Empirical data better support such a broader view and helps to comprehend the decision-making process more comprehensively. Among biological systems, dopamine-based reward-learning models have had a significant influence on the evolution of artificial intelligence. Artificial entities given reinforcement learning techniques based on dopaminergic systems have been able to learn by feedback and make decisions under uncertain conditions. The focus of these biologically-based methods on the translational applicability of dopamine studies demonstrates how neuroscience can inform the intelligent design of systems. However, despite these capabilities, there are serious challenges. Current paradigms often reduce complex brain interactions to simpler ones and do not account for the roles of other neuromodulator systems and higher-order cognition. Future studies must be directed toward developing more integrative models that describe the interaction among dopamine, cortical control systems, and the environmental context. A longitudinal, multimodal study will also help narrow model assumptions and improve predictive performance. In conclusion, dopamine reward learning models are among the fundamental approaches to the study of decision-making under ambiguity. Their further development and improvement are likely to yield greater knowledge about cognition, behaviour, and intelligent systems, thereby demonstrating their critical importance in both neuroscience and computer science.

7. Recommendations and Future Research

Based on the information and theoretical implications provided in this paper, several recommendations are offered to enhance the study of dopamine reward learning models and their use in decision-making under uncertainty.

- To begin with, scientific investigations of the future are advised to adopt more integrative modelling tools that can go beyond disdainful dopamine-based models. Although dopamine is critical for reward learning and decision-making, interrelations exist among neuromodulator systems, including serotonin, norepinephrine, acetylcholine, and others. Including these systems in computational models would yield more biologically realistic accounts of cognitive and

behavioural processes under uncertainty.

- Second, investigators should take up multimodal experimental approaches that integrate behavioural activities, neuroimaging and computer modelling. The application of techniques such as functional magnetic resonance imaging (fMRI), electroencephalography (EEG), and pharmacological treatment may provide complementary information on the evolution of dopamine signals in uncertain decision-making. Connecting brain inputs more directly to model parameters, such as learning rates and uncertainty sensitivity, will make dopamine-based models more explanatory and predictive.
- Third, the following research is to focus more on individual differences in dopamine activity. There is genetic variability, personality characteristics, and developmental effects on dopaminergic signalling and, consequently, decision-making behaviour. Modelling such individual-level variances can enhance insight into why individuals respond differently to uncertainty and risk. These methods are applied in individual interventions in both clinical and school settings. From a therapeutic perspective, neuropsychiatric and neurological diseases should be further investigated using dopamine reward-learning models.

8. Future Research

Further reports need to examine the mechanisms by which processes of disturbed dopamine cause maladaptive choices in interventions like addiction, Parkinson's disease and schizophrenia. Long-term studies on the evolution of dopamine signalling may be of interest for understanding disease progression and treatment efficacy. Computerised psychiatry measures can also be used to develop targeted treatment plans. In practice, the role of researchers and developers working in artificial intelligence should be to improve reinforcement learning algorithms based on the dopaminergic principle continually. Artificial agents in complex real-world scenarios can be enhanced by incorporating physiology-aware uncertainty models and adaptive learning schedules, thereby significantly improving their performance. Cooperation between neuroscientists and AI researchers is strongly recommended to ensure that computer models are grounded in empirical evidence. The ethical and societal implications of advances in dopamine-based decision models should be investigated further, particularly within artificial intelligence systems capable of influencing human behaviour. Insights into the impact of reward-based algorithms on decision-making could inform the more effective and efficient implementation of intelligent technologies. In conclusion, an interdisciplinary approach, methodological innovation, and greater emphasis on complexity, individuality, and real-world application are required to advance dopamine reward learning research.

9. References

- Attaallah, B., Petitot, P., Zambellas, R., Toniolo, S., Maio, M. R., Ganse-Dumrath, A., Irani, S. R., Manohar, S. G., & Husain, M. (2024). The role of the human hippocampus in decision-making under uncertainty. *Nature Human Behaviour*, 8(7), 1366-1382.
- Augustat, N., Endres, D., & Mueller, E. M. (2024). Uncertainty of treatment efficacy moderates placebo effects on reinforcement learning. *Scientific Reports*, 14(1), 14421.
- Ban, K. (2024). *On the computational and neural characterisation of reward learning behaviour* University of Glasgow].
- Ben-Zion, Z., & Levy, I. (2025). Representation of Anticipated Rewards and Punishments in the Human Brain. *Annual review of psychology*, 76(1), 197-226.
- Blanco-Pozo, M., Akam, T., & Walton, M. E. (2024). Dopamine-independent effect of rewards on choices through hidden-state inference. *Nature Neuroscience*, 27(2), 286-297.
- Bruckner, R., & Nassar, M. R. (2024). Decision-making under uncertainty. *PsyArXiv*.
- Cecchini, G., DePass, M., Baspinar, E., Andujar, M., Ramawat, S., Pani, P., Ferraina, S., Destexhe, A., Moreno-Bote, R., & Cos, I. (2024). Cognitive mechanisms of learning in sequential decision-making under uncertainty: an experimental and theoretical approach. *Frontiers in Behavioral Neuroscience*, 18, 1399394.
- Chou, K.-P., Hakimi, N., Hsu, T.-Y., & Smith, R. (2025). A Systematic Empirical Comparison of Active Inference and Reinforcement Learning Models in Accounting for Decision-Making Under Uncertainty. *Available at SSRN 5174669*.
- Dias Maile, A. A., Gründler, T. O., Fischer, A. G., Kurtenbach, H., Kaiser, L. F., Froböse, M. I., & Jocham, G. (2025). Bidirectional modulation of reward-guided decision making by dopamine. *Psychopharmacology*, 1-13.
- Dillon, D. G., Belleau, E. L., Origlio, J., McKee, M., Jahan, A., Meyer, A., Souther, M. K., Brunner, D., Kuhn, M., & Ang, Y. S. (2024). Using drift diffusion and RL models to disentangle effects of depression on decision-making vs. learning

in the probabilistic reward task. *Computational Psychiatry*, 8(1), 46.

- Feng, Y.-Y., Bromberg-Martin, E. S., & Monosov, I. E. (2024). Dorsal raphe neurons integrate the values of reward amount, delay, and uncertainty in multi-attribute decision-making. *Cell Reports*, 43(6).
- Gershman, S. J., Assad, J. A., Datta, S. R., Linderman, S. W., Sabatini, B. L., Uchida, N., & Wilbrecht, L. (2024). Explaining dopamine through prediction errors and beyond. *Nature Neuroscience*, 27(9), 1645-1655.
- Gershman, S. J., & Lak, A. (2025). Policy complexity suppresses dopamine responses. *Journal of Neuroscience*, 45(9).
- Ghosh, D., Ghosh, D., & Ghosh, D. (2025). Temporal Difference Learning and Dopaminergic Mechanisms in Bellman-Attention Transformers. In: Bellman-Attention.
- Kahnt, T., & Schoenbaum, G. (2025). The curious case of dopaminergic prediction errors and learning associative information beyond value. *Nature Reviews Neuroscience*, 26(3), 169-178.
- Khan, Z. (2025). Neurocognitive mechanisms of adaptive decision-making: An fMRI-based investigation of prefrontal cortex dynamics in uncertain environments. *Advances in Cognitive and Neural Studies*, 1(1), 20-27.
- Korbisch, C. C., & Ahmed, A. A. (2025). Rapid Dopaminergic Signatures in Movement: Reach Vigor Reflects Reward Prediction Error and Learned Expectation. *bioRxiv*, 2025.2003. 2024.645035.
- Kurtenbach, H., Froböse, M. I., Ort, E., Bahnert, B. H., Hirschmann, J., Butz, M., Schnitzler, A., & Jocham, G. (2024). A role for acetylcholine in reinforcement learning and decision making under uncertainty. *bioRxiv*, 2024.2009. 2020.614105.
- Liebana, S., Fritsche, M., & Lak, A. (2024). Dopaminergic computations for perceptual decisions. *Current Opinion in Behavioral Sciences*, 60, 101458.
- Mah, A., Golden, C. E., & Constantinople, C. M. (2024). Dopamine transients encode reward prediction errors independent of learning rates. *Cell Reports*, 43(10).
- Metha, J. A., Bertheau, M., Murawski, C., Hoyer, D., & Jacobson, L. H. (2025). Investigating the effects of Orexin receptor 1 antagonism on decision making under uncertainty. *Behavioural Brain Research*, 115691.
- Nussenbaum, K., & Hartley, C. A. (2024). Understanding the development of reward learning through the lens of meta-learning. *Nature Reviews Psychology*, 3(6), 424-438.
- Paunov, A., L'Hôtellier, M., Guo, D., He, Z., Yu, A., & Meyniel, F. (2024). Multiple and subject-specific roles of uncertainty in reward-guided decision-making. *bioRxiv*.
- Saxena, V. (2024). AI Meets the Brain: Integrating Cognitive and Behavioral Neuroscience Principles into Artificial Intelligence. Available at SSRN 5019283.
- Schultz, W. (2024). A dopamine mechanism for reward maximization. *Proceedings of the National Academy of Sciences*, 121(20), e2316658121.
- Skowron, A. (2024). *Neural variability as a marker of cognitive flexibility during learning and decision-making under uncertainty* Humboldt-Universität zu Berlin].
- Story, G., Kurth-Nelson, Z., Moutoussis, M., Igaya, K., Will, G.-J., Hauser, T., Blain, B., Vlaev, I., & Dolan, R. (2024). Discounting future reward in an uncertain world. *Decision*, 11(2), 255.
- Taira, M., & Sharpe, M. J. (2025). Complementary roles of serotonin and dopamine in model-based learning. *Current Opinion in Behavioral Sciences*, 61, 101464.
- Wang, Y., Lak, A., Manohar, S. G., & Bogacz, R. (2024). Dopamine encoding of novelty facilitates efficient uncertainty-driven exploration. *PLOS Computational Biology*, 20(4), e1011516.
- Wheeler, A.-R., Truettner, L. M., Boehnke, A., Kahanek, P., & Orsini, C. A. (2024). Sex differences in sensitivity to dopamine receptor manipulations of risk-based decision making in rats. *Neuropsychopharmacology*, 49(13), 1978-1988.