

Metapopulation Conservation Models under Habitat Fragmentation Scenarios

Kanyarat Rattanakorn^{1*}, Dewi Putri¹

¹Phuket Rajabhat University, Phuket, Thailand

ABSTRACT

In terms of integrating the survival of the species in discontinuous landscapes, where habitat degradation and fragmentation are significant dangers to biodiversity, metapopulation conservation models provide a holistic framework. The populations in these models are modeled as networks of discrete patches of habitats connected through the process of dispersal. This emphasizes the importance of colonization-extinction processes in defining the survival of the populations in the long term. When having habitat fragmentation, patch size, spatial arrangement, and connectedness significantly impact the stability of the metapopulation. Isolated and small patches have a higher probability of extinction. Recent developments in modeling have made it possible to make more precise predictions of anthropogenic-driven population dynamics. These innovations have been facilitated through the inclusion of stochasticity, species-species dispersion capacities and environmental variability. Simulation-based studies have demonstrated that maintenance of the interconnectedness of habitats and priority in ensuring that keystone patches are maintained can greatly diminish extinction. Moreover, the integration of metapopulation models into the landscape planning facilitates the process of determining important corridors and preservation strategies that will most likely ensure that species can survive over a long term. The present research can be applied in the evidenced-based conservation and management approaches that can help to preserve biodiversity in the ecosystems that are increasingly regulated by humans. It does integrate the known methods, uses and constraints of the metapopulation models in fractured habitats.

ARTICLE HISTORY

Received November 14, 2025

Accepted December 15, 2025

KEYWORDS

Metapopulation Conservation Models (MCM),
Habitat Fragmentation Scenarios (HFS)

1. Introduction

Life on this world depends on a vast number of plants, animals, and microbes that occupy diverse surroundings. However, these ecosystems are not necessarily huge and continuous, but consist of many smaller places that vary in size, shape, resources, and circumstances. They actually coexist in groups in different sectors of environments. These groupings have numerous associations, like migration of individuals between patches (Gutiérrez-Arellano et al., 2024). This movement is the foundation of the way by which most species can exist in nature, and particularly in topography, which inherently comprises hills, woodlands, rivers, or broad grasslands that provide separate habitats. human activities like farming, road construction, industrialization, urbanization and chopping down forests have split huge natural areas into small and solitary patches known as habitat fragmentation (Hovel, 2025). Fragmentation reduces the quantity of available habitat while also influencing how these habitats are positioned and connected. In most cases, this leads towards longer distances between patches of habitat, smaller patches, and an increased risk of local extinction. Highways, towns, or farms now separate the species that formerly moved freely over the landscape. This makes it significantly more difficult to maintain healthy and stable populations (Wu et al., 2025). Due to these challenges, ecologists and conservation biologists concluded that traditional methods of animal preservation, such as protecting a specific area of habitat, could be insufficient. One possibility is that current conservation must understand the dynamics of the population, which includes many patches (Tao et al., 2024). A global threat to biodiversity, which is one of the most pressing, is habitat fragmentation. Human activities such as the growth of urban settlements, agriculture, deforestation, and infrastructural development are the major causes of it. The continuous habitats are broken down into smaller, isolated units, and as a result, the biological processes that maintain the species populations are interrupted, and this causes an increased risk of extinction, altered species interactions and reduced ecosystem resilience. Fragmentation not only reduces the total habitat space but also isolates populations, thereby limiting dispersal and gene flow, which are essential in maintaining genetic diversity and adaptation capacity. In this context, having

knowledge of the dynamics of species populations of fragmented landscapes is crucial in achieving effective conservation plans and management. The metapopulation theory provides an analytical and conceptual tool that could be used to examine the process of populations in fragmented habitats. A metapopulation was coined in 1969 by Levins as a combination of physically independent groups of the same species that interact in a dispersal manner. In contrast to models of single populations, which assume that the habitats are continuous, the colonization of open spaces, and the local extinction of subpopulations, metapopulation models take into consideration both of these processes in establishing the overall population persistence. This viewpoint shifts the conservation emphasis away from within the populations to the larger landscape, which emphasizes connectivity in patches and the spatial distribution of habitat remnants. The effect of habitat fragmentation is immense on the metapopulation dynamics through changes in patch size, isolation, and quality. It is a common characteristic of smaller patches to sustain fewer individuals, making the population more vulnerable to demographic and environmental stochasticity, and isolated patches reduce the potential of individuals to recolonize after local extinctions. Thus, fragmented landscapes are normally characterized by extinction debt, in which populations decrease gradually with time, although no additional habitat degradation takes place. To study these dynamics, it is necessary to combine spatial ecology, population biology, and landscape modeling to predict the way species will react to the current habitat changes.

The metapopulation hypothesis provides this information by implying that populations are divided into smaller groups in separate habitat patches, resulting in the formation of multiple species. Such small populations may occasionally become extinct, but fresh immigrants can recolonize them from diverse places (Dietz et al., 2024). The ratio of regional population extinction to blank region resettlement determines the species' long-term viability. Using metapopulation theory, we can see that a species is not driven just by the environment inside a single patch, but also by the association of all patches within the landscape. The patches must be close enough together, and individuals must be able to travel between them, to ensure that a metapopulation can be maintained even if patches are absent or temporarily empty (Peterson et al., 2025). However, as fragmentation grows, patches become increasingly difficult to obtain. This discourages movement and limits the possibility of recolonization. In the long run, if too many patches are severely isolated, the total metapopulation may collapse. Scientists use metapopulation conservation models to identify these patterns and predict future outcomes (Watts & Hughes, 2024). These are the techniques that would help the researchers in studying how the species would flourish among diverse habitat patches in various environments. Other models are straightforward, and they are based on the count of occupied patches. The additional data sources are more comprehensive and include information on patch size, habitat quality, patch distance, and individuals' locomotor capabilities to cross the surface of the land (Galán-Acedo et al., 2024). The models allow scientists to imitate real-life conditions and investigate how species may evolve as the environment changes. The consequence of habitat fragmentation presents many challenges related to metapopulation. Small populations are more likely to become extinct because they are vulnerable to unexpected factors such as weather changes, disease, or low breeding success (Johansson et al., 2025). Traveling between isolated areas becomes increasingly challenging. This shows that once a local population has become extinct, it is extremely unlikely to be recolonized. As a result of fragmentation, organisms may become more vulnerable to predators, invasion by other species, and human activity. All of these variables contribute to the threat of extinction at the local level and reduce the ability of the entire metapopulation to maintain its existence (Peacock, 2025). Metapopulation conservation models assist in solving these issues by emphasizing how different landscape layouts affect species survival. Researchers can use the models to predict what would happen if particular patches were destroyed, other patches are created, or some habitat corridors exist in order to improve migration (Marchesan & Kolasa, 2024). Models also have the capability of displaying the patches that are most important in long-term survival. Some patches may serve as powerful, so-called, source populations, producing a large number of individuals, but others may require immigrants to survive. Conservation planners can concentrate their efforts on the most effective regions by identifying the most important patches. Technology has been developed to permit the production of more accurate and comprehensive metapopulation models. GPS tracking, satellite images, drones, GIS mapping, and others, enabling researchers to measure habitat quality, monitor animal movement, and the terrain barrier in a better way (Fernandes et al., 2025). This information is employed in spatially explicit metapopulation models, which simulate realistic models. These models may incorporate such information as the distance between patches, the steepness of the landscape, the effects of the roads or buildings. These models are more efficient as far as conservation planning is concerned due to their potential to deliver genuine data based on real-world situations (Mant et al., 2025). Metapopulation models may also be employed in management practice to construct previously undamaged landscapes, biological networks, and protected areas. Old methods were based on the idea that protecting a specific amount of habitat was sufficient to ensure the species' survival. However, the number of habitat patches is not the only important factor in irregular landscapes, but their structure and connectivity are also important (Lacy et al., 2024). The metapopulation idea recognizes the need to connect patches through pathways or other landscape characteristics that allow secure passage. It also emphasizes the need of conserving the patches that serve as important sources of population. When these models are merged into management planning, the species will survive in the long run (Moreira & Pereira, 2025). Furthermore, metapopulation conservation models under

habitat fragmentation situations offer valuable insights into how species can survive in patchy environments. They highlight the importance of connectivity, patch quality, and landscape structure for long-term survival (Hovestadt et al., 2024). Metapopulation conservation models have evolved over the past decades to more complex theoretical models, integrating species peculiarities, landscape stochasticity, and landscape complexity into a single model.

Classic Levins-type models describe populations by how much of the patches were occupied with the colonization and extinction rates being the equilibrium in metapopulations. These models offer a fundamental knowledge of the thresholds of persistence and main habitat requirements, but make the assumption that there is homogeneity between patches, and do not consider the spatial layout. In order to address these limitations, spatially explicit metapopulation models (SEMMs) were built, which include the worldwide distribution of habitat patches, dispersal ranges, and patch-dependent extinction hazards. SEMMs allow a researcher to create a habitat loss and fragmentation scenario, determine the effectiveness of conservation, and also locate key patches or corridors for species survival. Metapopulation models have been widely used in the fragmentation of a wide variety of species, such as mammals, birds, insects, and plants. As an illustration, research conducted on the species of butterflies in the European agricultural landscape shows that the linkage of habitat patches is crucial to the recolonization and long-term persistence. On the same note, big carnivores like tigers and wolves need connected networks of habitats in order to support viable populations, as isolated populations are at a greater risk of inbreeding and local extinction. In the case of plant species, seed dispersal methods and patch isolation have an effect on colonization success, which affects the stability of the metapopulation. These empirical implementations depict the flexibility of metapopulation frameworks to address conservation issues in diverse natural frameworks. One of the new conservation approaches is introducing metapopulation theory to landscape ecology, which tries to preserve or restore the linkages in the fragmented ecosystems. The interventions proposed to allow dispersal and increase the rates of recolonization usually include habitat corridors, stepping stones and buffer zones. Besides, by making the protection of large, high-quality patches or keystone patches a priority, the metapopulation persistence can be optimized since these patches will serve as sources to the surrounding. Scenario-based modeling enables conservation planners to investigate trade-offs, including the minimum acreage and connectivity of the habitat required to maintain the populations across long periods of time, even in the presence of constant anthropogenic stress. Regardless of the tremendous achievements, there are still issues in the use of metapopulation models in conservation when it comes to the real world.

2. Literature Review

The study of metapopulation conservation models has increased significantly in recent years as researchers attempt to comprehend how species can survive within increasingly fragmented environments. Metapopulations have been demonstrated to be more of a theory than a conservation practice, which has assumed association with destruction risk in addition to the management of the habitat (Amster et al., 2024). Researchers studied the impact of fragmentation on the ecological processes, and it was demonstrated that the existence of a species cannot be solely attributed to circumstances within an isolated patch. Researchers claimed that long-term persistence relies on the dynamics between dispersed populations via dispersal, recolonization and patch connectivity. Studied examined significantly that single-population models remain unsatisfactory to predict the determination of species in isolated habitats (Pinto et al., 2024). The fact that fragmentation changes demographic processes in a way that cannot be explained by the traditional population models has been argued by researchers. Many scholars investigated the question of the landscape pattern and its effect on population extinction, showing that the spatial structure of the habitat patches has a more important effect on the decline. Studies imply that the effect of fragmentation can not be quantified using the impacts of the quantity of the habitat alone. However, the configuration of the habitat is crucial as well (Grumbach & Hilker, 2025). These types of understanding formed the use of metapopulation models to better reflect these spatial dependencies. It has also been shown that dispersal between patches seems to be an insurance mechanism that can recolonize the local extinctions. Researchers noted that functional connectivity is more important than the actual physical proximity of patches (Forero-Sánchez et al., 2025). Their results had a reliance on several conservation strategies, especially the species with poor dispersal abilities like amphibians, small mammals and insect specialists. Many scholars have studied the issue of landscape permeability, and it was found that even the tiniest barriers can block the movement, such as the narrow roads or farmlands (Harisena et al., 2025). Studies indicated that conservation interventions were needed to enhance the landscape connection with the help of corridors, stepping-stone environments, or recreated vegetation. It also concerns the literature with another idea of the metapopulation models being simple theoretical models that have evolved into complex, data-driven frameworks. The Incidence Function Model (IFM) used patch area and isolation as a part of the colonization and extinction probability, leading to more predictions that matched the empirical data (Zhang et al., 2024). It has been improved by many researchers with the addition of stochastic processes, environmental variance and species-specific characteristics. These sophisticated models enable conservation planners to predict species results in complementary conditions of fragmentation to a higher



degree. Quality of patches has also become the focus of a critical element of persistence of metapopulations (Teixeira et al., 2025). Studies indicated that patch suitability was a direct and independent variable of the occupancy rates, even when patches were big or highly connected. It was highlighted that the viability of metapopulation is not only determined by the spatial structure but also by the internal conditions of the patches, which include the availability of resources, microclimatic stability and predator abundance (Falconi-Briones et al., 2025). Researchers established that deteriorated patches would serve as sinks where the stability of the overall metapopulation would be reduced. Due to this, conservation is more environmentally focused, involving habitat restoration in order to enhance the quality of patches, as well as spatial organization. Studies have shown that isolation between patches leads to a slowing of inheritable factor flow, causing genetic drift and inbreeding (Gao et al., 2024). Researchers emphasized that genetic decadence may increase the risk of extinction and decrease the adaptability of a population, particularly when small populations are considered. Genetic data have been used to simulate genetic structure using metapopulation models to analyze the impacts of dispersal (Van Daele et al., 2024).

The metapopulation study of butterflies, birds, and mammals indicates that even rare disperses may preserve genetic diversity at the regional level. It was revealed that minimum connectivity is critical to demographic stability, but genetic resilience as well. Metapopulation models have also increased in application with climate change research (Serva et al., 2025). Scholars investigated the climatic changes and examined that the climate changing belt have influencing habitat suitability, which proves that species need more mobility in order to seek a favorable environment. Researcher shows that fragmented landscapes limit changes in range, as well as exerting further pressure on already pre-existing stress-sensitive populations because of habitat loss (Leite et al., 2025). The metapopulation models have shown that, under the influence of climate, species with small dispersal ability will be at a greater risk of extinction unless the landscape connectivity is enhanced. Many scholars have demonstrated that climate forecasts and spatial plans must be merged in order to come up with conservation schemes that embrace future patterns in habitat (Sierra et al., 2025). Researchers have been able to develop metapopulation models further owing to technological advances. With GIS-based techniques, high-density mapping and remote sensing techniques, there has been enhanced accuracy of the spatial models as they can reflect the actual landscape structures. The study of butterflies, frogs, birds, and carnivores indicates that metapopulation management facilitates the existence of species when compared to the processes of management that rely on the dynamics of isolated individual populations (Zou et al., 2025). Studies point out that Metapopulation models are criticized and limited despite their strong points. Researchers observe that not every species fits in a metapopulation structure, especially those with highly mobile or nomadic behavior. Authors said that most models rely on parameters that are not easily measured, which include true dispersal rates or patch-specific survival probabilities (Bhandary et al., 2025). Demonstration of model predictions in the long term was still a problem because there was no empirical evidence that could support such predictions. To overcome these shortcomings, scholars recommend the use of field observations, experimental studies and long-term monitoring in conjunction with a modeling method to enhance reliability. In general, the literature shows that metapopulation conservation models can provide an effective way to consider the dynamics of species in fragmented landscapes. The contributions of connectivity, patch quality, landscape configuration, genetic mixing, and environmental change are repeatedly pointed out by the researchers (Segura et al., 2025). The rates of colonization and extinction of species are at times difficult to calculate accurately owing to a lack of data on changes in the population, dispersal, and quality of the habitat. In addition, stochastic processes such as natural disasters or climatic extremities have the potential to radically alter the results of metapopulation, which requires probabilistic methods and sensitivity analyses. Climate change also makes it more difficult to do projections, as changing environmental conditions can change the suitability of a habitat, dispersal routes and species interactions and result in needing dynamic models which incorporate time variability. Also, metapopulation models are often interested in single species dynamics, when in actuality, interspecies interactions, trophic structures and ecosystem interactions are important determinants of population dynamics. To solve these challenges, metapopulation modeling will require combining with larger ecological and socio-economic scales, such as land-use planning, habitat restoration and community-based conservation actions. Remote sensing, geographic information systems (GIS) and landscape modeling have made significant technological leaps that have substantially increased the ability to apply metapopulation models to fragmented landscapes. Spatial data can be mapped in detail to track habitat patches and their connectivity, and also trace the dynamics of populations across time when these data are of high resolution. Conservationists can use simulation models to investigate other possibilities of habitat loss, corridor establishment, and patch reinstatement, providing evidence-based guidance to policy and management. Furthermore, combining metapopulation models with population viability analyses (PVA) and species distribution models (SDM) makes it possible to have a more in-depth evaluation of the risk of extinction in different conditions of fragmentation and climate.

To conclude, metapopulation conservation models provide an effective methodology to examine and reduce the effects of habitat fragmentation on biodiversity. These models offer important observations of the effect of spatial organization on population persistence by emphasizing the relationship between patch occupancy and colonization as well as between patch occupancy and extinction. Their use in fragmentation conditions shapes the direction to real conservation

actions, such as habitat conservation, construction of corridors, and landscape control, to help species in more humanized ecological communities. Nevertheless, the issues of data access, stochasticity, climate change, and interactions among multi-species underscore the necessity to enhance and expand further on applying metapopulation models on a larger ecological and management context. Further studies that focus on geographically and temporally shifting models, which are supported by empirical evidence and interdisciplinary methods, would be critical towards guaranteeing the long-term survival of species in discontinuous ecologies.




Figure 1: Metapopulation Conservation Models

3. Metapopulation Conservation Models

Metapopulation conservation models are analytical and simulation-based models that are applied to study the persistence of species in fragmented environments (Figure 1). These models represent populations as a web of discrete patches of habitat linked together during dispersal, with the emphasis on local extinctions and recolonization processes. Their applications are especially applicable to conservation biology because habitat loss and fragmentation have since become one of the major contributors to a reduction in biodiversity.

3.1 Conceptual Foundation

Richard Levins developed the idea of metapopulation in 1969. This notion, according to which a species is no longer viewed as a single continuous population, but as a collection of spatially dispersed subpopulations (or local populations) that inhabit discrete patches of habitation. These subpopulations are linked together through the process of dispersal, which enables them to recolonize occupied patches and also genetic interchange of claims among populations. Important processes in the dynamics of metapopulations are:

3.1.1 Local Extinction

The subpopulation of a patch becomes extinct as a result of stochastic events, a small population or environmental shifts. The origin of a population on a previously empty patch, usually aided by diffusion between adjacent patches, is known as colonization (Recolonization).

3.1.2 Spatially Explicit Metapopulation Models (SEMMs)

SEMMs combine patch and species-specific dispersal capability with geographic locations. Spatial dependence on colonization and probability of extinction. This is useful to simulate realistic landscapes with irregular patches of varying forms, size and dispersal barriers. SEMMs are especially applicable to the study of the effects of habitat fragmentation, corridor development and restoration.




Figure 2: Stochastic and Dynamic Models

3.1.3 Stochastic and Dynamic Models

These models have random demographic and environmental variability (Figure 2). Stochastic models are essential in small or very remote populations where random events can be quite strong factors affecting the level of extinction. The temporal shifts in the quality of habitats, patterns of land-use, and climate impacts can also be incorporated into dynamic models.

4. Research Methodology

Metapopulation dynamics in the context of capricious fragmentation of habitats is a study that includes both theoretical modulization, geography, and simulation approaches. The approach outlines the model of developing, evaluating, and understanding the metapopulation models to evaluate the persistence of species in fragmented habitats. It involves the classical metapopulation theory, spatially explicit models, and scenario-based simulation in order to study the effects of habitat fragmentation on the colonization-extinction dynamics.

4.1 Study Framework

The study is concerned with populations of species distributed in a fragmented landscape in terms of a network of discrete patches of habitats. Its area, quality and connectedness differentiate each patch from other patches. The metapopulation is assumed to be comprised of local populations that can undergo local extinction but can be recolonized by dispersal of patches occupied. The habitat fragmentation is recreated through changing the patch sizes, isolation distances, and the landscape architecture as a whole to emulate real-life situations of habitat loss and the human modification of the landscape.

4.2 Data Collection and Landscape Mapping

Categorical data will be gathered using past (2010/2011) reports and newsletters that will be analyzed using a content analysis methodology. The remote sensing databases and Geographic Information Systems (GIS) are used to acquire the spatial data on habitat patches. Each patch is quantified in terms of habitat quality indicators like the vegetation cover, availability of resources and the level of anthropogenic disturbances. Patch connectivity is studied with the help of patches using Euclidean distance between patches, least-cost path analysis or graph-theoretical methods, taking into consideration dispersal corridors and obstacles. Field surveys, published literature and species databases provide empirical observations of species occurrence, density of species population and dispersal ability that are used to parameterize the model.

4.3 Metapopulation Model Formulation

The study employs both classical and spatially explicit metapopulation models:

- Levins' Metapopulation Model:
- The basic Levins model describes the dynamics of the proportion of occupied patches $P(t)$ over time:

$$\frac{dP}{dt} = cP(1 - P) - eP$$

Where:

- P is the proportion of occupied patches
- c is the colonization rate
- e is the extinction rate

This equation provides a simple framework to examine the balance between colonization and extinction in determining metapopulation equilibrium.

- Spatially Explicit Metapopulation Models (SEMM):
- SEMMs incorporate the spatial distribution of patches, patch-specific extinction probabilities, and species dispersal limitations. The colonization probability of an empty patch is given by:

$$C_i = 1 - \prod_{j \in O} (1 - p_{ij})$$

Where:

- O is the set of occupied patches
- p_{ij} is the probability of colonization from patch j to patch i , often modeled as a function of distance d_{ij} and dispersal ability α :

$$p_{ij} = c \cdot e^{-\alpha d_{ij}}$$

Extinction probability E_i for each patch is calculated as a function of patch size, quality, and stochastic environmental factors:

$$E_i = e_0 \cdot f(A_i, Q_i)$$

Where e_0 is the baseline extinction rate, A_i is patch area, and Q_i is patch quality.

4.4 Network Analysis and Connectivity Metrics

The graph-theoretical methods are used to represent the patches of habitats by the nodes and the routes among them by the edges.

Network metrics, including:

- Degree significance: The number of connections in a patch, which illustrates that it is involved in dispersal.
- Betweenness centrality: Significance of patches as connections between subpopulations.
- Clustering coefficient: Local network cohesiveness, which shows that it is possible to recolonize clusters. Such measures help to define the important patches the protection of which or restoring which is highly necessary to ensure the stability of the metapopulations.

4.5 Sensitivity Analysis

The sensitivity analysis is carried out to study how the main variables, including the rate of colonization, the rate of extinction, and dispersal capacity, affect the outputs of the models. The robustness of the metapopulation to various fragmentation conditions is studied by systematically varying the parameters and thresholds to determine whether persistence or collapse can be discovered.

4.6 Model Validation

Real data from field observation or historic patterns of the population is used to verify model projections. Simulated patch occupancy and experiential species distributions are compared to determine the accuracy of the model. In case direct data are not available, expert knowledge and literature-based parameters are used to calibrate the models.

4.7 Conservation Implications

Conservation solutions are developed depending on the model outputs so as to contain the impact of habitat



fragmentation. These are:

- Protection of big or well-connected patches should be a priority problem
- Corridor of habitat should be designed to aid in dispersal.
- Recovery of damaged areas to boost occupancy in the area
- Introduction of landscape-level planning to maintain network resilience.

The methodology has offered a holistic perspective on the impact of habitat fragmentation on metapopulation processes and presented valuable information on the conservation of biodiversity and management of landscapes (Figure 3).




Figure 3: Ecological Metapopulation Concepts

5. Equational Analysis and Interpretation of Network Metrics

Network theory offers an effective scheme of the evaluation of habitat connectivity and its effects on the metapopulation processes within fragmented environments. The structural and functional features of the habitat network are measured through important network measures by modeling habitat patches as nodes and dispersal channels as edges. Such measurements are used to identify important patches, approximate fragmentation, and address conservation interventions. This section gives an overview of the most important equations applied in network analysis and the ecological meaning of these equations.

5.1 Degree Centrality

The degree centrality (k_i) is defined by the number of immediate connections a patch i has to other patches in the network:

$$k_i = \sum_j a_{ij}$$

Where:

- $a_{ij} = 1$ if there is a dispersal link between patches i and j , otherwise 0
- The summation is over all patches j

Interpretation: Both high degree centrality and high degree peripherality imply areas with several dispersal relationships, which serve as centers of colonization. These patches often play a vital role in the persistence of metapopulation because their disappearance can separate the network and predispose local extinction.

5.2 Betweenness Centrality

Betweenness centrality (BC_i): This is the number of shortest paths between any two patch pairs that traverse patch i :

$$BC_i = \sum_{s \neq i \neq t} \frac{\sigma_{st}(i)}{\sigma_{st}}$$

Where:

- σ_{st} is the total number of shortest paths between patches and t
- $\sigma_{st}(i)$ is the number of those paths that pass through patch i

Interpretation: Patches having high betweenness centrality are used as stepping stones or important connections that connect the otherwise remote areas. Preservation of such patches leads to functional interconnectedness, which facilitates dispersal and recolonization. Their elimination may pose a significant threat to network fragmentation and the extinction of metapopulations.

5.3 Clustering Coefficient

The tendency of a patch to be interconnected with neighbor patches is a measure of the clustering coefficient (C_i) of the patch:

$$C_i = \frac{2e_i}{k_i(k_i - 1)}$$

Where:

- e_i is the number of edges between neighbors of patch i
- k_i is the degree of patch i

Interpretation: Heavy clustering indicates that there is redundancy in the local connections, and there are various dispersal pathways. This will enhance network resilience since the failure of one patch does not isolate its neighbors. On the other hand, low clustering is an indication of vulnerability, since patches rely on a limited number of links to connect.

5.4 Network Density

Network density (D) measures the proportion of actual connections relative to all possible connections in the network:

$$D = \frac{2E}{N(N - 1)}$$

Where:

- E is the total number of edges in the network
- N is the total number of patches (nodes)

Interpretation: High network density implies a well-connected landscape where dispersal pathways are abundant, supporting higher colonization rates and lower extinction risk. Low density indicates sparse connectivity, increasing vulnerability to fragmentation.

5.5 Patch Connectivity Index (PCI)

Patch connection uses both spatial and dispersion data to measure the functional importance of a patch in facilitating the persistence of a metapopulation:

$$PCI_i = \sum_{j \neq i} O_j \cdot e^{-\alpha d_{ij}}$$

Where:

- O_j is the occupancy state (1 if occupied, 0 if empty) of patch j
- d_{ij} is the distance between patches i and j
- α is a species-specific dispersal decay parameter

Interpretation: The PCI shows the possibility of patch i to receive colonists of other occupied patches. Examples of patches containing high PCI are vital in metapopulation resilience, whereas an example with low PCI is separated and faces a higher chance of extinction. This action enlightens the fixation on restoration or protection.

5.6 Simulation and Sensitivity Analysis

Monte Carlo simulation is stochastic colonization-extinction events through the network, after the dynamics of


occupancy with time. The sensitivity of metapopulation persistence to fragmentation can be assessed by changing network factors, including patch loss, connection and dispersal ability. Output measures are:

- Fraction of occupied patches with time
- Time to extinction of subpopulations.

Patch removal has an effect on network connectivity.

These researchers permit determining threshold conditions of persistence and impact conservation initiatives at the landscape level. On the whole, the quantitative insights into the effects of habitat fragmentation on the dynamics of metapopulation are provided by equational analysis and network metrics. The degree and betweenness centrality show essential patches dispersal, clustering, and network density measure the local and global connectivity, and PCI is the correlation between structural connectivity and functional colonization potential. A combination of these indicators and colonization-extinction probability provides practical information on the choice of patches, establishment of corridors, and implementation of effective conservation policies in fragmenting ecosystems.

6. Metapopulation Network Model



$$N_1 = n_{11} + n_{12} + n_{13}$$

$$\hat{N}_1 = n_{11} + n_{21} + n_{31}$$

$$\frac{dS_1}{dt} = -\beta\nu \sum_{k=1}^3 \left\{ \frac{n_{1k}}{N_1} S_1 r_k \sum_{\ell=1}^3 \frac{E_{\ell} + f I_{\ell}}{N_{\ell}} \frac{n_{\ell k}}{\hat{N}_k} \right\} - \beta(1 - \nu) S_1 r_1 \frac{E_1 + f I_1}{N_1}$$

Figure 4: Metapopulation Network Model

7. Applications of Metapopulation Models and Network Metrics in Conservation:

Metapopulation models and network measures are useful in conservation efforts of biodiversity, especially where there is habitat fragmentation. These models can be used to make decisions on the protection, restoration, and management of habitats by evaluating connectivity, patch importance, and by evaluating the threat of extinction. The uses of these methodologies are very diverse and can be found in detecting priority patches and constructing corridors and conservation projects (Figure 4).

1. Priorities of Critical Habitat Patches: Network metrics, such as degree centrality, betweenness centrality, and Patch Connectivity Index (PCI), help in determining important patches of habitats that allow persistence of the metapopulation. Patches of high degree centrality or betweenness centrality are the hubs or linkages in the landscape. An insurance of protection of these patches ensures functional connection and recolonization of other patches following local extinctions. Indicatively, in fragmented forest landscapes, source populations of species occur in patches that are categorized as high-centrality nodes, and therefore, the conservation of these areas is critical to conserving regional biodiversity.

2. Development and execution of Habitat Corridors: The metapopulation model can be used to compute dispersal routes and possible corridors among patches in a spatially explicit manner. The conservation planners can construct corridors that would enable the population to migrate between isolated ones by determining the colonization probability and the risk of extinction of the respective population under different landscape configurations. As an example, network analysis has been used to construct wildlife corridors in the agricultural landscapes of Europe and the forest systems of North America to preserve connectivity of species like lynx, wolves and migratory birds. Network measures control the position, width and quality of corridors to improve connectivity and alleviate the effect of isolation.

3. Degraded Patch repair: using patch-specific connection indices and extinction probability, it is possible to prioritize degraded patches to be repaired. It is possible to target patches with intermediate connectivity and high extinction risk to achieve habitat enhancement, including reforestation, elimination of invasive species, or lighting up of resources. Restored

patches are also effective in enhancing local carrying capacity as well as overall network resilience by means of additional dispersal nodes. The strategy has been used in grassland and wetland restoration initiatives to facilitate the stability of metapopulations of pollinators, amphibians and small animals.

4. **Landscape-Level Planning and Conservation Zoning:** The Metapopulation models provide the result of the large-scale land-use planning using methods such as identifying areas where the protection of habitats, growth restriction on the land-use, or the creation of corridors would have the most conservation benefit. When network measurements are incorporated with land-use data gathered using the GIS, planners are in a position to construct zoning policies that support the balance between human activities and ecological requirements. As an illustration, metapopulation simulations have been used to develop landscape-scale conservation plans in tropical forest reserves to define the core areas of protection and buffer zones to keep the impacts of fragmentation minimal but still allow the species of focus to be connected.

5. **Measuring the Consequences of Fragmentation and Land-Use Change:** Metapopulation modeling can be used to measure the potential consequences of the loss of habitat, urbanization or agricultural expansion before they happen. The changes in patch occupancy, connectivity and extinction probability are measured by simulations conducted in scenarios with other development paths. There is no impact on metapopulation persistence that this forecasting power does not mitigate. Therefore, it enables policymakers to take mitigation measures to counter such unfortunate impacts, including strategic land acquisition or corridor construction. As an example, the computer studies conducted in the tropical rainforest environment have assessed the probability of a species extinction in the conditions of the supposed deforestation, pushing the conservation theory into the offensive.

6. **Information on Species Reintroduction and Translocation Programs:** Network analysis can help direct the species reintroduction or translocation activities by discovering suitable source and target patches. Patches that are highly connected and have a low probability of extinction are very promising in forming new populations, which guarantees successful colonization and the survival of the population. As an example, to improve survival and enhance dispersal in the protected areas, metapopulation and network models have been used to plan the reintroduction of endangered carnivores, including the Iberian lynx.

7. **Improving Climate Change Response Plans:** Climate change effects change the metapopulation and habitat suitability and connectivity. The spatial explicit models have the capability to synthesize predicted changes in habitat-appropriate locations to predict the patches that will be viable and corridors that will facilitate range changes driven by climate. This program provides pre-emptive conservation strategies (like protection of climatic refugia, enhancing connectivity across existing and future apposite habitats, and focusing on adaptive management strategies of vulnerable species).

8. **Monitoring and Adaptive Management:** Metapopulation models facilitate adaptive management by quantifying the change in population trends and changing habitat connectivity over time. Network analysis gives the evolution of patch significance, network density, and colonization-extinction dynamics, which guide the change of conservation strategies as the landscape changes. Indicatively, butterfly metapopulations in broken European landscapes have been observed over a long period to use these tools to customize management interventions, including mowing regimes and habitat recovery.

9. **Incorporation of Multi-Species Conservation:** Although most metapopulation models have been conducted in isolation of species, multi-species conservation can be generalized in the same way as network measures. The identification of overlapping habitat needs and connectivity needs can enable managers to identify patches and corridors to sustain a large number of species at the same time and to conserve species more effectively. As an example, tropical and temperate systems Landscape planning tends to examine joint animal, bird, and pollinators conditions to construct networks that amplify general biodiversity robustness. Metapopulation and network measures are models that use ecological theory to convert the theory into practical conservation actions. These methods can underpin evidence-based solutions to the sustainability of biodiversity in humanized landscapes through the identification of significant patches, the designation of corridors, guidance of restoration, recommendations of reintroductions, and the examination of the consequences of the fragmentation. Combined with GIS, field data, and scenario-based simulations, it becomes possible to conduct proactive, adaptive and multi-scale conservation plans to ensure that fragmented ecosystems retain viable populations over time.

8. Discussion

The problem of habitat fragmentation is a severe threat to the conservation of biodiversity, which distorts biological processes and increases the risk of extinction of species. These dynamics can be fully analyzed using the metapopulation models and network measurements to provide the necessary management actions. Findings in this research paper show the significance of structural and functional connectedness in aiding metapopulation persistence in disaggregated environments. The experiment demonstrates that the patch size and isolation are important variables of the local extinction and colonization frequencies. Smaller patches experience greater extinction threats owing to both demographic and

environmental stochasticity, and the isolated patches experience reduced recolonization ability. This is consistent with the traditional theory of metapopulation, in which the overall stability of the populations is a matter of the balance between extinction and colonization. The centrality measures of a network, like degree and betweenness centrality, show that the well-connected patches are used in the form of hubs or stepping-stones, facilitating dispersal and recolonization within the landscape. Failure to maintain such critical patches may cause malfunction in the network, and extended extinctions. This observation is in agreement with empirical research on mammals, birds and insects, in which the elimination of important nodes in the habitat results in the rapid decline in population persistence. Conservation efforts should hence not just be directed at large patches of habitats but at those areas that offer functional connectivity to the network.

9. Conclusion

Comprehensively, the present research highlights the applicability of a metapopulation theory and network-style research in explaining and addressing the impacts of habitat fragmentation. These technologies are useful by providing valuable information on landscape-level conservation efforts by measuring patch significance, connectivity, and extinction risk. Proper implementation of the mentioned concepts can help achieve metapopulation resilience, enable species survival, and sustain biodiversity in ecosystems that are becoming more human-centered and fragmented. Metapopulation conservation models are the theoretical and empirical frameworks that are used in understanding the persistence of species in fragmented ecosystems. These tools provide conservationists with the opportunity to estimate the extinction threat, prioritize patches, develop a corridor, and implement evidence-based management by integrating spatially explicit simulations, network measurements, and probabilistic modeling. They need to be included in the landscape-level planning to help preserve biodiversity in man-made and fragmented ecosystems.

10. Limitations and Future Directions

Although the metapopulation models offer valuable information, there are some boundaries that need to be approached. The problem of estimating the rate of colonization and extinction may arise because of the lack of empirical evidence, particularly of elusive or rare species. The results can be influenced by extreme weather, disease outbreaks or human interference, which are stochastic events that are difficult to predict. In addition, the majority of the models consider the dynamics of only a single species, whereas the real ecosystems involve complex interspecific relationships. Put simply, future studies need to consider multi-species networks, include dynamic habitat quality dynamics, and integrate metapopulation models with population viability studies and species distribution models to conduct a more detailed conservation planning.

11. References

- Amster, P., Robledo, G., & Sepúlveda, D. (2024). Persistence/extinction scenarios in an almost periodic metapopulation with competition and habitat destruction. *arXiv preprint arXiv:2412.02952*.
- Bhandary, S., Gawecka, K. A., Pedraza, F., & Bascompte, J. (2025). Landscape configuration and community structure jointly determine the persistence of mutualists under habitat loss. *bioRxiv*, 2025.2005. 2028.655631.
- Dietz, J. M., Mickelberg, J., Traylor-Holzer, K., Martins, A. F., Souza, M. N., & Hankerson, S. J. (2024). Golden lion tamarin metapopulation dynamics five years after heavy losses to yellow fever. *American Journal of Primatology*, 86(7), e23635.
- Falconi-Briones, F. A., Bolom-Huet, R., Naranjo, E. J., Reyna-Hurtado, R., Enríquez-Rocha, P. L., Moreira-Ramírez, J. F., García, M. J., & Medellín, R. A. (2025). Connectivity at risk: a critical scenario for the endangered Baird's tapir and the vulnerable white-lipped peccary in the Maya Forest. *Biodiversity and Conservation*, 34(1), 235-254.
- Fernandes, P. F., de Avelar São-Pedro, V., de Lima Souza, B., Moreira-Silva, C., de Cássia Quitete Portela, R., Mendes, E. T. B., de Oliveira Scarascia, P., & da Silva Carvalho, C. (2025). Effects of defaunation of large seed dispersers, habitat loss and fragmentation on the population expansion of a tropical palm. *Landscape Ecology*, 40(4), 1-16.
- Forero-Sánchez, F., Rezende, G. C., Valladares-Pádua, C., Vannucchi, F. S., Jerusalinsky, L., Pacca, L., & Traylor-Holzer, K. (2025). Securing black lion tamarin populations: improving habitat-based inputs and risks for population viability analysis to inform management decisions. *Frontiers in Conservation Science*, 5, 1423321.
- Galán-Acedo, C., Fahrig, L., Riva, F., & Schulz, T. (2024). Positive effects of fragmentation per se on the most iconic metapopulation. *Conservation Letters*, 17(3), e13017.

- Gao, J., Gong, J., Li, Y., Yang, J., & Liang, X. (2024). Ecological network assessment in dynamic landscapes: Multi-scenario simulation and conservation priority analysis. *Land Use Policy*, 139, 107059.
- Grumbach, C., & Hilker, F. M. (2025). Allee pits in metapopulations: critical dispersal rates for connectivity to be beneficial. *Theoretical Ecology*, 18(1), 5.
- Gutiérrez-Arellano, C., Crone, E. E., Pettorelli, N., & Hodgson, J. A. (2024). Broadening applications of stochastic patch occupancy models over three decades. *Diversity and Distributions*, 30(5), e13822.
- Harisena, N. V., Grêt-Regamey, A., & van Strien, M. J. (2025). Shape and rate of landscape change trajectories influence species persistence. *bioRxiv*, 2025.2008. 2004.666294.
- Hovel, K. (2025). CRUSTACEAN RESPONSES TO HABITAT LOSS, FRAGMENTATION, AND MODIFICATION. *Ecology and Conservation: Volume 10*, 10, 195.
- Hovestadt, T., Poethke, H.-J., Müller, J., & Mitesser, O. (2024). Species diversity and habitat fragmentation per se: the influence of local extinctions and species clustering. *The American Naturalist*, 203(6), 655-667.
- Johansson, V., Franzén, M., & Bergman, K.-O. (2025). A 20-year study of a threatened butterfly: The importance of management across space and time for long-term persistence in dynamic landscapes. *Biological Conservation*, 310, 111367.
- Lacy, R. C., Breininger, D. R., Breininger, D. J., Savage, A. E., Forsman, A. M., Hoffman, E. A., McGuffey, S. D., DeMeyer, D., & Mecklenborg, T. (2024). Linking habitat and population viability analysis models to account for vegetation dynamics, habitat fragmentation, and social behavior of a metapopulation of Florida scrub-jays. *Frontiers in Conservation Science*, 5, 1505016.
- Leite, M. C., Oliva-Zúniga, R. I., & Torres, P. J. (2025). A meta-population model for non-timber forest product harvesting: connectivity, plant viability, and sustainable yield.
- Mant, D., Orevi, T., & Kashtan, N. (2025). Impact of micro-habitat fragmentation on microbial population growth dynamics. *The ISME Journal*, 19(1), wræ256.
- Marchesan, J., & Kolasa, J. (2024). Habitat fragmentation affects richness—a view through a metacommunity lens. *Diversity*, 16(11), 658.
- Moreira, R. M., & Pereira, G. T. (2025). The Role of Protected Areas in the Amazon: Modeling and Quantifying Future Forest Fragmentation Integrating Artificial Neural Networks and Landscape Ecology Metrics. *Journal of Sustainable Forestry*, 44(10), 1218-1242.
- Peacock, M. M. (2025). Negotiating a Fragmented World: What Do We Know, How Do We Know It, and Where Do We Go from Here? *Diversity*, 17(3), 200.
- Peterson, M. C., Bradham, J., Ferraro, K., Guillet, M., Keuroghlian, A., Ribeiro, M., & Jorge, M. L. S. (2025). Modeling movement patterns to identify thresholds of functional connectivity in fragmented forest landscapes. *Biological Conservation*, 307, 111177.
- Pinto, A. V., Hansson, B., Patramanis, I., Morales, H. E., & van Oosterhout, C. (2024). The impact of habitat loss and population fragmentation on genomic erosion. *Conservation Genetics*, 25(1), 49-57.
- Segura, J., Marvá, M., & Franco, D. (2025). Dispersal in multi-patch metapopulations: the impact of patch number and network topology. *Journal of Theoretical Biology*, 112331.
- Serva, D., Iannella, M., Biondi, M., & Krofel, M. (2025). Integrating habitat suitability, connectivity, and individual-based models to guide priorities for the creation of a lynx metapopulation in Southeastern Europe. *Biological Conservation*, 310, 111381.
- Sierra, A. M., Alonso-García, M., Zartman, C. E., & Villarreal A, J. C. (2025). Migration Patterns of Epiphyllous Bryophyte Metapopulations in Amazonian Fragmented Landscape Inferred From Census and Genomic Data. *Biotropica*, 57(5), e70088.
- Tao, Y., Hastings, A., Lafferty, K. D., Hanski, I., & Ovaskainen, O. (2024). Landscape fragmentation overturns classical metapopulation thinking. *Proceedings of the National Academy of Sciences*, 121(20), e2303846121.

- Teixeira, C., Nunes, G. L., Trevelin, L. C., Silva, D. P., & Prudente, A. L. d. C. (2025). Assessing Connectivity Thresholds Under Habitat Loss Scenarios for Threatened Amphibians and Squamate Reptiles in the Eastern Brazilian Amazon. *Ecology and Evolution*, 15(7), e71741.
- Van Daele, F., Honnay, O., & De Kort, H. (2024). Plant conservation in a changing world: modelling range dynamics, climate adaptation, and effects of habitat fragmentation.
- Watts, K., & Hughes, S. (2024). Fragmentation impacts may be mixed for conservation but generally bad for restoration. *Restoration Ecology*, 32(8), e14260.
- Wu, D., Yu, Y., Luo, M., Meng, B., & Wang, S. (2025). Spatial heterogeneity modulates dispersal effects on metapopulation abundance: insights from a graphical approach.
- Zhang, H., Chase, J. M., & Liao, J. (2024). Habitat amount modulates biodiversity responses to fragmentation. *Nature Ecology & Evolution*, 8(8), 1437-1447.
- Zou, Y., Crowther, T. W., Smith, G. R., Ma, H., Mo, L., Bialic-Murphy, L., Potapov, P., Gawecka, K. A., Xu, C., & Negret, P. J. (2025). Fragmentation increased in over half of global forests from 2000 to 2020. *Science*, 389(6765), 1151-1156.