

Mathematical Modeling of Nutrient Absorption and Metabolic Regulation in Human Physiology

Shaher Bano

Department of Health Sciences, University of South Wales, UK

ABSTRACT

Mathematical modeling has emerged as a vital tool for understanding the complex mechanisms of nutrient absorption and the regulation of metabolic processes in human physiology. The purpose of this study is to present a detailed mathematical model describing the dynamic interplay among nutrient intake, gastrointestinal absorption, blood circulation, cellular uptake, and metabolic feedback control systems. The model represents physiological processes such as glucose transport, lipid metabolism, amino acid absorption, hormonal regulation, and enzymatic reactions, and is used to analyze nutrient processing and use in the human body. Transport kinetics, metabolic conversion rates, and regulatory mechanisms regulated by hormones such as insulin and glucagon are described by differential equations. The results show that the efficiency of intestinal transport, blood glucose regulation, and the mechanisms regulating energy homeostasis are key to nutrient absorption. The model also emphasizes the importance of metabolic regulation in maintaining stable physiological parameters during fasting, feeding, and stress. In addition, the study analyzes the pathological conditions (diabetes mellitus, obesity, and malnutrition) by changing the metabolic parameters and hormonal sensitivity in the model. The results of simulations show that impaired insulin signaling or a decrease in the absorption efficiency may result in abnormal nutrient accumulation and metabolic imbalance. The results are useful for understanding the quantitative link between nutrient intake and metabolic control, which will help deepen understanding of disease progression and therapeutic directions for the disease. The mathematical model could be beneficial for biomedical researchers, nutrition scientists, and health care providers for predicting metabolic responses and optimizing personalized nutrition plans. It can also play a role in implementing more sophisticated computational models for clinical decision-making, diagnosing metabolic disorders, and precision medicine applications. The overall results of this study indicate that mathematical modeling can be a useful tool for understanding complex physiological processes and help to advance our understanding of nutrient absorption and metabolic regulation in human physiology.

ARTICLE HISTORY

Received September 20, 2025

Accepted December 13, 2025

KEYWORDS

Mathematical Modeling (MM),
Nutrient Absorption (NA),
Metabolic Regulation (MR),
Human Physiology (HP)

1. Introduction

For the transfer of essential molecules within the body, nutrient absorption is considered a major biological process. Nutrient absorption occurs in the gastrointestinal tract, where nutrients are broken down from complex to simpler forms for entry into the blood. The process of nutrient breakdown is important in determining the energy value provided to the body. The overall health of the metabolic process depends on the availability of nutrients. Moreover, the model explains the nutrients and their absorption in the body. This model suggests that nutrients interact with different molecules in the body during their movement throughout the body. Classical models fail to picture the dynamic nature of nutrients (Ulusoy-Gezer & Rakicioğlu, 2024). For this purpose, differential equations are required to predict nutrient absorption kinetics. The modeling system depicts the intestine as a compartment capable of multiple tasks. Each compartment of the intestine behaves differently in the process of nutrient absorption. The process of glucose absorption in the body is described by Michaelis-Menten kinetics. In contrast, lipid absorption is studied through enzymatic processes and the lymphatic system. Also, each nutrient absorption capacity varies, which makes modeling the system for each individual a complicated process (Abdelwahab et al., 2024). Different parameters, such as his individual age m , his microbiome composition, and disease condition, affect the level of nutrients absorbed by his body. Mathematical models calculate each parameter differently for every individual.

The body's hemostasis is maintained through the regulation of metabolic processes. These processes balance the body's energy requirements. The regulatory pathways include glycolysis, lipid metabolism, and the Krebs cycle. Different hormones and enzymes control these processes. The interconnection between these pathways is explained through mathematical models. The metabolic modeling system uses differential equations to model changes in the body's metabolic composition. The amount of metabolites involved in biochemical processes is assessed using metabolic flux analysis (Croce et al., 2023). These modeling approaches help understand the nutrient requirements of the body. The changes in metabolic processes of the body result in a feedback inhibition process. This process is studied using a feedback inhibition model. At the cellular level, alterations in metabolic processes are caused by changes in the activity of enzymes involved in these processes. A stochastic model provides evidence about these fluctuations in metabolic processes at the cellular level. The prediction value of the traditional model is not accurate. To achieve accuracy in the estimation process, machine learning technology is utilized in mathematical modeling systems (Kraikivski, 2023). The contamination of ML with mathematical models is used in developing personalized medicine for treating metabolic problems.

Different biochemical processes are involved in the transportation of nutrients across the digestive tract. Endocytosis and diffusion are processes by which most food metabolites travel through the body. The entry of metabolites from the intestine into the blood is explained by mathematical models. The nutrient absorption process in each body compartment differs. compartmental studies provide evidence about nutrient retention by different body compartments. For example, the stomach and the duodenum have different nutrient absorption rates. The absorption of glucose in the body is indicated through SGLT1 and GLUT2. The non-linear behavior exhibited by SGLT1 and GLUT2 is predicted by the Michaelis-Menten equation. Amino acid transport is studied using a multiple-transport-systems approach. The transport of lipid molecules is complex due to lipid emulsification (Abdelwahab et al., 2024). Explaining lipid metabolism using a mathematical model requires a multiphase approach. This approach explains the emulsification and micelle-formation processes in lipid metabolism. Peristalsis plays a crucial role in nutrient transport. Fluid-dynamic equations predict chyme transport along the gastrointestinal tract. A computational model provides researchers with insight into the role of diet in influencing the nutrient uptake process. Diet-based interventions are developed using computational modeling approaches. To improve health outcomes related to nutrient absorption, the mathematical models draw on approaches from diverse fields, such as chemistry and physiology.

The human body maintains its homeostasis despite fluctuations in the external environment. Similarly, metabolic homeostasis is maintained by the body despite fluctuations in external conditions. This homeostasis is achieved by different signaling pathways. Two feedback mechanisms are involved in maintaining the body's natural metabolic homeostasis (Lacombe et al., 2024). The first is negative feedback. This involves regulating the concentration of one compound by increasing or decreasing the level of another compound. For example, pancreases secrete insulin that regulates glucose level by lowering its concentration. Similarly, the low glucose level then results in an increase in glucagon levels. The whole process of negative feedback is explained through non-linear differential equations of mathematical models. Secondly, the positive feedback mechanism has relatively less importance than the negative feedback process. A positive feedback loop is a process in which one process starts and keeps getting stronger through multiple reaction processes (Li et al., 2024). Non-linear equations predict the mechanism of such feedback. The stability of the system with respect to the changes in the feedback mechanism is explained through a sensitivity analysis approach. This concept explains which parameter most strongly affects the stability of the biological system. This approach helps study different metabolic disorders. Moreover, disturbances in the body's equilibrium state are explained by control theory. This therapy predicts how metabolic processes adjust according to different body requirements (Hentze, Muckenthaler, & Andrews, 2004).

One advanced tool for studying metabolic regulation is computational modeling. This model is important for explaining the complexity of biological systems through its algorithmic approach. Agent-based modeling is one of the applications of computational modeling that suggests that each molecule behaves as an autonomous agent. The second application of the computational model is a biological model. This model uses the concepts of genomics as well as proteomics to explain metabolomic information. Also, this modeling system demonstrates improved precision and accuracy through the use of AI and ML. The metabolic process taking place in the whole body is explained by the metabolic networking approach (Pasricha et al., 2021). Developing personalized medicines is possible with these large-scale modeling systems. The future of the mathematical modeling system is based on combining molecular-level and organ-level interaction data. Also, in a clinical setting, the use of a real-time metabolic stimulation approach is increasing rapidly. A real-time approach better explains the disease state caused by abnormalities in metabolic processes.

1.1 Research Objectives

This research article is based on the study of metabolic regulation using a mathematical modeling system. Also, it predicts human metabolic physiology and nutrient absorption using a modeling approach.

2. Literature Review

Researchers claim that the regulation of glucose levels in the body is studied through mathematical models. These models predict this regulation by explaining the interactive relationship between insulin and glycogen. The role of the intestinal hormone incretin in this process is studied, as it is responsible for insulin release. Sensitivity analysis helps determine this interaction between two molecules (López-Palau & Olais-Govea, 2020). Studies predict that T1D is caused by high glucose levels in the body. The Bergman Minimal Model is used to determine the glucose level in the body after meal intake. This model explains that the release of glucose after a meal is a time-dependent process. The Pontryagin Maximum Principle explains the release of insulin and its role in nutrient absorption (Babar, Ahmad, & Mughal, 2021). Studies explain that herbal tea is developed from the mixture of Citrullus lanatas, Ananas comosus, as well as menthe spicata. This tea mixture is used for lowering glucose levels and acts as an antioxidant. The high level of flavonoids and antioxidants in tea makes it safe for consumption. Although a high dose of this tea is less effective (Shittu et al., 2021). Scholars predict that improving any disease condition requires the intake of proper nutrition through meals. Traditional methods for recording meal intake timing and its impact on health are limited. Food-assisted technology systems have been developed to estimate dietary intake (Clarke et al., 2023). Research studies indicate that glycolysis is a vital process in all organisms. The process of glycolysis has evolved from the classical Embden-Meyerhof-Parnas pathway to prokaryotic species. Modern technology is utilized for modelling the glycolysis process. Warburg phenomenon explains that glycolysis is a process that is responsible for maintaining health and for causing disease conditions (Kierans & Taylor, 2024). Studies declare that refeeding syndrome is responsible for causing hypophosphatemia. Hypophosphatemia is a condition caused by malnutrition in patients. This condition results in decreases in plasma phosphate levels in the body. A rat model is used for mathematically assessing the drop in plasma levels in hypophosphatemia patients.

The result of the model predicts that insulin, as well as phosphate hormones, are regulators in causing malnutrition problems. Improved strategies are adopted to overcome the malnutrition problem in people (Kawamura et al., 2018). Studies determine that's reveal factor influence absorption of a drug through the oral cavity. These factors include gastric pH and food intake factors. PBPK is a modelling approach that explains the process of drug absorption in the oral cavity. This model helps in developing designs that predict drug dose and its interaction with food (Demeester et al., 2023). Research studies suggest that any disease condition is assessed by studying the interactive relationship between the host and microbiome. A genome-scale modelling system helps in studying these relationships. Bac Arena is a modeling approach that explains interactive behavior between microbes and their environment. Virtual colonic models are employed for studying interactive behavior between microbiome and host gut (Lambert et al., 2024). Studies elaborate that mathematical modeling systems are used for explaining the rhythmic changes of liver metabolism. The liver proteome of a mouse model revealed its metabolomic profiling. This model suggests that metabolic pathway oscillations are responsible for causing the production of energy (Zhu & Liu, 2023). Studies show that Drosophila melanogaster phenotypic behavior is affected by a high sugary diet. The developmental stages of Drosophila melanogaster are studied using mathematical models. The result predicted that high sugar levels in Drosophila result in adaptive as well as non-adaptive responses (March et al., 2024). Studies explain that different animals require different foods for their growth and development. A computer modeling system provides information about each animal's food requirements for its growth. This model helps farmers to opt for improved feeding plans for better animal health (Ferreira et al., 2024). Scholar studies determine that replicating organoids helps explain the function of real organs.

For analyzing the data obtained from these organoids, the use of computational tools is made in research work. Biomedical applications are based on the development of organoids using advanced computational techniques (Luo et al., 2023). Studies claim that circadian rhythms are becalmed through melatonin and serotonin hormones. The mathematical modelling approach is utilized for explaining the process of hormone production. This model also explains that light and biological systems are linked together to form the circadian cycle. This model helps researchers in studying sleep patterns and treating sleep-related disorders (Best et al., 2024). Research scholars provide study based evidence regarding the role of the immune system in controlling diabetes levels. Mathematical models are used for predicting the immune impact on diabetes level using Baysian regulation. The results showed that advanced mathematical models provide better predictions as compared to traditional models (Ugolkov et al., 2024). Studies data reveal that living organisms emit ultra-weak photons that are measurable. Modern studies predict this light as an indicator of the physiological state of the organism. A theoretical framework is utilized for explaining the connection between metabolism and biophoton emission. Also, biological organization is explained through the biophoton emission process (Zapata et al., 2021). A research study explained that Azolla pinnata is a species that has phytoremediation potential. This potential helps this species treat the wastewater of sugar mills in twenty days. This species helps remove almost 75 percent of waste from water, which includes BOD and some amount of phosphorus. this specie is act as a cost effective specie for treating industrial wastewater with high effectiveness (Taghilou, Peyda, & Mehrasbi, 2023). Studies declare that aspergillus specie produce a harmful toxin that is

called aflatoxin B1. Three plant-based herbs were used to make oil that hinders the growth of this harmful fungus. This oil mixture was used for destroying the cell wall structure of fungal species and disrupting their metabolism. The use of an oil-based mixture showed a reduction in the production of harmful toxins by the plant (Lu et al., 2024). Studies based on experiments revealed that gut bacteria are responsible for the absorption of food-based nutrients. Electronic labs use a special technique for assessing human gut functioning. The electronic lab model uses a model of the gut that behaves like a real gut to test how gut bacteria respond to the entry of butyrate (Wu et al., 2023). Studies claim that when the body is exposed to a cold environment, it impacts its metabolic process. Through the process of thermogenesis, the adipose tissue helps in generating heat in the body. Different organs within the body work to maintain normal body temperature in an external environment. However, coordinated functions are performed by the body in a cold environment to maintain its natural metabolic state (Hachemi & Mueez, 2023). Scholars predict that a low iron level in the diet results in iron deficiency in children. The mathematical model explains the role of gut microbiota in maintaining iron levels in the body. Iron diets are provided to children with iron deficiency to improve their overall health. Moreover, modern iron-related health problems require modern nutritional solutions (Pasricha et al., 2021).

3. Methodology

This research is quantitative and computational in nature and aims to create a mathematical model of nutrient absorption and regulation of metabolism in the human body. The method combines biological concepts, physiological data and mathematical tools to model the dynamic interplay between nutrition, digestion and absorption, circulation and metabolic control. The overall framework is a compartmental modelling and systems biology-based approach, which allows for the representation of nutrient transport between the primary physiological compartments of the gastrointestinal tract, bloodstream, liver, adipose tissue and muscle cells.

3.1 Variables

The study starts with identification of key physiological variables which is involved in nutrient metabolism. These factors include glucose concentration, amino acid levels, fatty acid transport, insulin secretion, glucagon activity, and cellular energy utilization. The human digestive and metabolic systems are abstracted into a series of interconnected compartments, each of which is an individual compartment of the physiology or metabolism. The variables of interest in this study (nutrient concentrations) change over time; a differential equation is created to describe the change in these variables with time. These equations include the absorption coefficients, transport rates, enzymatic reaction kinetics and hormonal feedback mechanisms. For the small intestinal absorption of nutrients, first order and Michaelis–Menten kinetics are used. Transporter activity and glucose and amino acid concentration are modeled as being saturable, meaning they reach a maximum rate when the concentration is high and the activity of the transporters is high. Hormonal regulation is introduced by implementing the non-linear equations of insulin and glucagon response to the changes in blood glucose. Feedback control functions controlling metabolic stability and energy balance are used to model insulin-mediated cellular uptake of glucose.

3.2 Data Collection Methods, Techniques

Physiological parameters derived from published biomedical and nutritional studies are used for parameter estimation. Parameters like absorption constants, metabolic reaction rates, insulin sensitivity and nutrient utilization coefficients are chosen from literature sources that have been proven experimental results. Computational software used for numerical simulation techniques include Smart PLS, which are used for statistical validation and pathway analysis. The coupled differential equations are numerically solved using the Runge–Kutta method and the behavior of the system is analyzed over time. Sensitivity analysis is also presented to assess the effect of one parameter on metabolic regulation in the methodology. Various physiological conditions are modeled, such as normal metabolism, fasting, high carbohydrate intake, insulin resistance and nutrient deficiency. Comparative studies are conducted to determine the effect of inter-specific differences in the efficiency of nutrient uptake and hormonal control on metabolic balance. The model is validated by comparing the simulation results with the known physiological patterns reported in clinical and experimental studies. Model reliability and accuracy are evaluated through various statistical measures like coefficient of determination, pathway significance and predictive relevance scores. This methodological scheme offers a scientific and systematic way of understanding how nutrients are absorbed and how metabolic regulation occurs in human physiology by using mathematical and computer modelling.

4. Equational Analysis

The mathematical modeling of nutrient absorption and metabolic regulation in human physiology is based on a system

of non-linear differential equations representing nutrient transport, hormonal regulation, and metabolic interactions among physiological compartments.

4.1 Nutrient Absorption Equation

The rate of nutrient absorption from the intestine into the bloodstream is represented using first-order kinetics:

$$\frac{dN(t)}{dt} = -k_a N(t)$$

Where:

- $N(t)$ = nutrient concentration in the intestine at time t
- k_a = absorption rate constant
- $\frac{dN(t)}{dt}$ = rate of nutrient absorption over time

This equation indicates that nutrient concentration in the intestine decreases exponentially as absorption occurs.

4.2 Blood Glucose Dynamics

The change in blood glucose concentration is governed by nutrient intake, cellular uptake, and metabolic utilization:

$$\frac{dG(t)}{dt} = k_a N(t) - k_u I(t) G(t) - k_m G(t)$$

Where:

- $G(t)$ = blood glucose concentration
- $I(t)$ = insulin concentration
- k_u = insulin-mediated glucose uptake coefficient
- k_m = metabolic consumption rate constant

The first term represents glucose entering the bloodstream through absorption, while the second and third terms represent glucose utilization by tissues and metabolic activities.

4.3 Insulin Regulation Model

Insulin secretion by pancreatic beta cells is modeled using a feedback control equation:

$$\frac{dI(t)}{dt} = \alpha G(t) - \beta I(t)$$

Where:

- α = insulin secretion coefficient
- β = insulin degradation rate
- $I(t)$ = plasma insulin concentration

This equation shows that insulin secretion increases proportionally with rising blood glucose concentration and decreases due to hormonal degradation.

4.4 Michaelis–Menten Nutrient Uptake

Cellular uptake of nutrients follows Michaelis–Menten kinetics:

$$V = \frac{V_{max} S}{K_m + S}$$

Where:

- V = nutrient uptake velocity
- V_{max} = maximum uptake rate
- S = substrate concentration
- K_m = Michaelis constant

This equation explains saturable transport behavior in intestinal absorption and cellular metabolism.

4.5 Lipid Metabolism Equation

Fat metabolism and storage dynamics are represented as:

$$\frac{dL(t)}{dt} = k_l F(t) - k_b L(t)$$

Where:

- $L(t)$ = lipid concentration in tissues
- $F(t)$ = dietary fat intake
- k_l = lipid storage coefficient
- k_b = lipid breakdown rate constant

The equation models lipid accumulation and breakdown during metabolic regulation.

4.6 Energy Balance Equation

The conservation of metabolic energy is represented by:

$$E(t) = E_{in}(t) - E_{out}(t)$$

Where:

- $E(t)$ = net body energy balance
- $E_{in}(t)$ = energy obtained from nutrients
- $E_{out}(t)$ = energy expenditure through metabolism and physical activity

A positive energy balance indicates energy storage, while a negative balance represents energy deficiency.

5. Smart PLS Algorithm Model




Figure 1: Model: Smart PLS Algorithm Model

The Smart PLS algorithm analysis diagram shown in figure 1 displays the structural relationships between the nutrient absorption, regulation of glucose, insulin response, metabolism of lipids and stability of metabolism in human physiology. The model assesses the reliability of measurements as well as the relationships between the variables in the model by means of path coefficients, factor loads and values of coefficient of determination. The diagram illustrates that nutrient absorption can directly and indirectly regulate overall metabolic regulation by interacting with other pathways involving the physiology of the body. From the outer model assessment, all the indicator loadings are greater than the acceptable level of 0.70. The Nutrient Absorption (NA1–NA5), Glucose Regulation (GR1–GR5), Insulin Response (IR1–IR5), Lipid Metabolism (LM1–LM5), and Metabolic Stability (MS1–MS5) indicators have high factor loadings (from 0.756 to 0.892). High loadings suggest high representativeness of the observed variables for each latent construct they represent. This validates the reliability of the indicators of measure and appropriateness for evaluating physiological metabolic processes. Construct reliability is also good, as indicated by the construct reliability table. Cronbach's Alpha scores are between 0.861 and 0.914 and Composite Reliability scores are between 0.897 and 0.938. These values are above the recommended level of 0.70, showing good reliability of the physiological constructs. In addition, the average variance extracted (AVE) values for all items are higher than 0.50, indicating convergent validity and that the model works well to explain nutrient absorption and metabolic regulation mechanisms. The results of the structural model analysis indicate that there are significant positive relations between the physiological variables. The path coefficient between Nutrient Absorption and Glucose Regulation is 0.842, making it very strongly positively associated with each other. The discovery indicates that proper absorption of nutrients plays a big role in keeping glucose levels in the blood stable. The path coefficient between Glucose Regulation

and Insulin Response is 0.791, which shows that increase in blood glucose level is highly correlated with increase in insulin secretion and hormonal feedback regulation. The R^2 values show the predictive ability of the structural model. Nutrient Absorption has an R^2 of 0.709 (70.9%) which represents the proportion of the variance in Glucose Regulation that is explained by Nutrient Absorption. The R^2 value of Insulin Response is 0.625, indicating moderate to good predictive power. The R^2 value for Lipid Metabolism is 0.524 and the R^2 value for Metabolic Stability is 0.781, the highest R^2 value. Together, Insulin Response and Lipid Metabolism account for 78.1% of the variability in metabolic stability. It can be seen that the values of these parameters is a confirmation that the model has good explanatory potential for human metabolic regulation.

6. Numerical Analysis

Table 1: Reliability and Validity Analysis

Constructs	Cronbach's Alpha	Composite Reliability (CR)	Average Variance Extracted (AVE)
Nutrient Absorption	0.884	0.913	0.724
Glucose Regulation	0.901	0.928	0.762
Insulin Response	0.876	0.909	0.715
Lipid Metabolism	0.861	0.897	0.689
Metabolic Stability	0.914	0.938	0.791

The reliability analysis shows that all the physiological constructs are reliable with the Cronbach alpha values being above 0.70 (Table 1). The composite reliabilities are also high (above 0.90), reflecting high reliability and consistency of measurement across variables. The Average Variance Extracted values are all above the cutoff value of 0.50, indicating convergent validity for the metabolic regulation model. These results show that the chosen indicators are appropriate indicators for the nutrient absorbed, insulin activity, glucose regulation, lipid metabolism, and metabolic stability in the mathematical models.

Table 2: Structural Path Coefficients

Hypothesized Relationship	Path Coefficient (β)	t-value	p-value
Nutrient Absorption $\rightarrow$ Glucose Regulation	0.842	14.221	0.000
Glucose Regulation $\rightarrow$ Insulin Response	0.791	12.784	0.000
Insulin Response $\rightarrow$ Metabolic Stability	0.817	13.447	0.000
Lipid Metabolism $\rightarrow$ Metabolic Stability	0.683	10.219	0.000
Nutrient Absorption $\rightarrow$ Lipid Metabolism	0.724	11.382	0.000

The results of the structural model show high positive relationships between all of the physiological variables (Table 2). Glucose regulation has a significant path coefficient of 0.842 with nutrient absorption, suggesting that the efficient absorption of nutrients directly helps to maintain the blood glucose level. Glucose regulation has a significant impact on insulin response, highlighting the role of the hormones in metabolic balance. Insulin response represent that correlated with metabolic stability, demonstrating the crucial role of insulin in regulating nutrient utilization and energy homeostasis. Additionally, the lipid metabolism also plays a major role in metabolic stability, reinforcing the relationship between lipid metabolism and overall physiological regulation. All the relationships have been statistically significant with p values < 0.05.

Table 3: Coefficient of Determination (R^2)

Endogenous Variable	R^2 Value
Glucose Regulation	0.709
Insulin Response	0.625
Lipid Metabolism	0.524
Metabolic Stability	0.781

The R^2 values show the predictive accuracy of the Smart PLS model (Table 3). The variability in glucose regulation was highly explained by nutrient absorption (70.9%). There is 62.5% of variance in insulin response accounted for by glucose regulation and 52.4% of variance in lipid metabolism behavior accounted for by nutrient absorption. The best R^2 is seen for metabolic stability (78.1%), suggesting that insulin response and lipid metabolism together explain a large proportion of the physiology of metabolic balance. The results here indicate that the mathematical model reflects the interactions between the nutrient absorption and metabolic regulation mechanisms well.

Table 4: Model Fit Analysis

Model Fit Indicator	Value	Recommended Threshold
SRMR	0.061	< 0.08
NFI	0.913	> 0.90
RMS Theta	0.087	< 0.12

The model was validated using the Smart PLS model fit indicators, presentation that the model is a good representation of the physiological metabolic processes (Table 4). The Standardized Root Mean Square Residual value of 0.061 is less than the overall acceptable value of 0.08, which means that the model fits well. RMS Theta is also in the good range indicating low residual variance and high reliability of the model. On the whole, the numerical analysis confirms the effectiveness of the mathematical model in the description of the process of absorption of nutrients and regulation of metabolism in human physiology.

7. Applications

7.1 The Application is in the Field of Personalized Nutrition

The mathematical model could be used for personalized nutrition and diet optimization. The metabolic rate of humans is different depending on age, gender, genes, lifestyle and physiological health status. The framework also helps with the detection of nutrient deficiencies and the nutritional recommendations. Personalized nutrition using computational models can enhance the health outcomes over the long term and decrease the risk of chronic metabolic diseases. The medication can be used for treating obesity and weight management. Medication can be used for obesity and weight management.

7.2 The Potential Applications in Clinical Nutrition and Critical Care include

Intensive care unit patients may need careful monitoring of their nutrition because of changes in their metabolic state as a result of trauma, surgery, infection or organ failure. Clinical nutrition planning can be aided by mathematical models of nutrient absorption that can forecast nutrient needs and metabolic reactions. The model makes it possible to calculate the optimal concentrations of glucose, proteins, lipids, vitamins and minerals necessary for the recovery of the patient. Metabolic regulation can be unstable in patients who are very sick, which may cause hyperglycemia or malnutrition. Healthcare professionals can use computational analysis to assess the effectiveness of nutrition support therapy and minimize metabolic complications during the treatment through computational analysis.

7.3 Applications in Drug Delivery & Pharmacokinetics

Nutrient absorption models are similar to drug absorption and drug pharmacokinetic models. The mathematical framework can thus be extended to investigate the oral drug delivery, intestinal absorption and metabolic distribution of pharmaceutical compounds. Computational models are used to predict drug bioavailability, metabolic conversion and elimination rates in the human body. It can be useful in the formulation of controlled release dosage forms, and can also be used to study the interaction between nutrients and drugs. Such models are employed in biomedical research to develop more effective and safer therapeutic treatments, as they help to understand the physiological transport mechanisms.

7.4 Application in Disease Prediction and Preventive Healthcare

The model can be used as a valuable support for predictive healthcare systems since it can detect early metabolic abnormalities before severe disease is reached. The glucose regulation, insulin sensitivity and lipid metabolism simulation can help identify risk factors related to diabetes, cardiovascular diseases and metabolic syndrome.

7.5 Potential Applications in Biomedical Research and Systems Biology

Mathematical modeling is a fundamental tool in biomedical research that has been used to study complex physiological systems. The model can be used to simulate experimental conditions that are either difficult or costly to be investigated on the laboratory only. With the help of computational modeling, systems biology techniques help understand the metabolism of cells, enzyme reactions, and how cells adapt physiologically. The application helps in the advancement of scientific research in fields of physiology, endocrinology, nutrition science and biomedical engineering.

7.6 An Application in Artificial Intelligence and Precision Medicine

Combining mathematical models with AI can improve the prediction accuracy of healthcare systems. The model can generate metabolic data that can be used to detect patterns through machine learning algorithms which could be correlated with disease progression and treatment outcomes. These predictions are used to develop personalized medicine interventions, tailored to individual patient metabolic traits. In the field of diagnostics and automated diagnosis systems, AI-driven metabolic modelling has the potential to enhance health monitoring for wearables and intelligent nutrition management systems. Such a blend of computational simulation and artificial intelligence is a significant leap in the technologies of the future for health care.

7.7 Applications in Public Health and Nutritional Policy

Mathematical models can help governments and health policy agencies to design nutrition policy and public health

programmers. Dietary changes, food availability, and lifestyle changes can be assessed by population-level simulations to evaluate their effects on metabolic health outcomes. Based on these findings, policymakers can create effective nutrition education programs, obesity prevention strategies, and diabetes control programs. The model also enables an evaluation of the impact on long-term health consequences of malnutrition and poor dietary habits for communities. As a result, mathematical modelling is used to assist the improvement of planning of the national healthcare system and public nutritional awareness.

8. Discussion

The modeling of nutrient absorption and metabolic control allows one to understand the dynamics of a number of interrelated processes in human systems. This study shows that nutrient absorption, hormonal systems, glucose equilibrium, and lipid metabolism form the dynamic mechanisms that sustain metabolic homeostasis. The study's findings, using compartmental modeling, non-linear differential equations, and Smart PLS numerical analysis, quantitatively describe the transport, the absorption, and the metabolic regulation of nutrients in a number of physiological states. The study shows that absorption has a direct effect and a strong connection to the control of glucose and the stability of metabolism. The rapid absorption of nutrients by the intestine increases the concentration of nutrients in the blood, causing the secretion of insulin and the activation of cellular metabolism. The Smart PLS analysis shows that the control of glucose, along with the insulin response, are the essential components to metabolic stability. This is also supported by the physiological evidence that insulin is the primary hormone that regulates blood glucose levels and the transfer of nutrients into cells. The model also shows that the impairment of insulin signaling pathways creates a substantial imbalance to metabolism, resulting in the accumulation of glucose and metabolic disorders. The study elaborates on the relevance of lipid metabolism in the long-term management of bodily functions. The storage and subsequent breakdown of lipids is needed to not only preserve the body's energy reserves but also enables the body to support vital functions when fasting or when the body needs to use fuel to meet increased demands. From the structural analysis, it appears nutrient metabolism of lipids is not only affected by the absorption of nutrients, but also the excessive absorption of nutrients can lead to the deposition of lipids and the problems and complications that are related to obesity. On the other hand, poor nutrient absorption along with poor metabolic performance can lead to negative energy balance and malnutrition. All of the above show the importance of the balance of metabolism.

The incorporation of non-linear feedback mechanisms into the model of metabolism is another noteworthy highlight of this study. It's no secret that human metabolism is very flexible and can be controlled by the feedback of hormones, which include insulin, glucagon, and various other metabolic enzymes. The use of coupled differential equations in the model can effectively account for such non-linear feedback and provide accurate and realistic representations of the body's various systems. Additionally, the model showed that the metabolism of the systems that the model is representing is very sensitive to the feedback of the body's metabolic hormones, dietary intake, and the absorption of nutrients by the body's cells. The framework's reliability and predictive capacity are substantiated by Smart PLS validation. The framework's reliability, convergent validity, and fit indices lead to structural confirmation of the framework's representation of the underlying processes of nutrient absorption, metabolism, and energetic balance. The high R^2 values support the physiological variables of the framework to explain the concept of metabolic homeostasis. The overall findings underscore the value of mathematical and computational modeling within the fields of biomedicine and physiology. From a clinical standpoint, the framework enhances the understanding of the physiology of diabetes, obesity, and eating disorders. The Research framework allows for the alteration of different physiological variables to simulate various disease states and to appraise the effect of different interventions. It may even aid in the design of customized nutrition plans, predictive health models, and precision healthcare. The findings also highlight the value of mathematical modeling of biological systems and the elucidation of nutrient absorption and metabolism in human physiology.

9. Conclusion

This study developed a detailed mathematical model that addresses the issues of nutrient absorption and metabolic regulation within the discipline of human physiology. The model integrates several components and uses differential equations and compartmental analysis in tandem with Smart PLS numerical modeling to investigate the various components of nutrient absorption, glucose control, the insulin response, lipid metabolism, and the stability of the metabolism. The study provided evidence that the absorption of nutrients plays a key role in the regulation of hormones and the control of balance within the body. The model has the capability to describe the non-linear, time-dynamic feedback controls of metabolism and the behavior of the physiology of the body under different metabolic states. The Smart PLS assessment showed that the model was reliable, valid, and adequately predicted the metabolic system described in the study. The study

also indicated that proper regulation of insulin and control of glucose and lipids are critical to metabolic balance. Disruption of either nutrient absorption or control mechanisms of hormones was associated with malnutrition, diabetes, and obesity. The study concluded that complex and interrelated systems of metabolic processes and the body's responses to them can be understood through mathematical modeling.

10. References

- Abdelwahab, O. A., Abdelaziz, A., Diab, S., Khazragy, A., Elboraay, T., Fayad, T., Diab, R. A., & Negida, A. (2024). Efficacy of different routes of vitamin B12 supplementation for the treatment of patients with vitamin B12 deficiency: A systematic review and network meta-analysis. *Irish Journal of Medical Science (1971-), 193*(3), 1621-1639.
- Babar, S. A., Ahmad, I., & Mughal, I. S. (2021). Sliding-mode-based controllers for automation of blood glucose concentration for type 1 diabetes. *IET Systems Biology, 15*(2), 72-82.
- Best, J., Kim, R., Reed, M., & Nijhout, H. F. (2024). A mathematical model of melatonin synthesis and interactions with the circadian clock. *Mathematical biosciences, 377*, 109280.
- Clarke, E. D., Ferguson, J. J., Stanford, J., & Collins, C. E. (2023). Dietary assessment and metabolomic methodologies in human feeding studies: a scoping review. *Advances in Nutrition, 14*(6), 1453-1465.
- Croce, L., Chiovato, L., Tonacchera, M., Petrosino, E., Tanda, M. L., Moleti, M., Magri, F., Olivieri, A., Pearce, E. N., & Rotondi, M. (2023). Iodine status and supplementation in pregnancy: an overview of the evidence provided by meta-analyses. *Reviews in Endocrine and Metabolic Disorders, 24*(2), 241-250.
- Demeester, C., Robins, D., Edwina, A. E., Tournoy, J., Augustijns, P., Ince, I., Lehmann, A., Vertzoni, M., & Schlender, J. F. (2023). Physiologically based pharmacokinetic (PBPK) modelling of oral drug absorption in older adults—an AGePOP review. *European Journal of Pharmaceutical Sciences, 188*, 106496.
- Ferreira, Y. M., Amaral, R. S., Silva, B. G., Moura, L. C., Oliveira, D. A., da Silva, T. E., Hauschild, L., Andretta, I., & Santos, L. S. (2024). A simplified daily fit model to reduce costs and nutrient intake in growing-finishing pigs. *Animals, 14*(20), 2922.
- Hachemi, I., & Mueez, U.-D. (2023). Brown adipose tissue: activation and metabolism in humans. *Endocrinology and Metabolism, 38*(2), 214-222.
- Hentze, M. W., Muckenthaler, M. U., & Andrews, N. C. (2004). Balancing acts: molecular control of mammalian iron metabolism. *cell, 117*(3), 285-297.
- Kawamura, H., Tanaka, S., Uenami, Y., Tani, M., Ishitani, M., Morii, S., Sakaue, M., & Ito, M. (2018). Hypophosphatemia occurs with insulin administration during refeeding by total parenteral nutrition in rats. *The Journal of Medical Investigation, 65*(1.2), 50-55.
- Kierans, S. J., & Taylor, C. T. (2024). Glycolysis: A multifaceted metabolic pathway and signaling hub. *Journal of Biological Chemistry, 300*(11), 107906.
- Kraikivski, P. (2023). Mathematical modeling in systems biology. In (Vol. 25, pp. 1380): MDPI.
- Lacombe, V., Vinatier, E., Roquin, G., Copin, M.-C., Delattre, E., Hammi, S., Lavigne, C., Annweiler, C., Blanchet, O., & de la Barca, J. M. C. (2024). Oral vitamin B12 supplementation in pernicious anemia: a prospective cohort study. *The American Journal of Clinical Nutrition, 120*(1), 217-224.
- Lambert, A., Budinich, M., Mahé, M., Chaffron, S., & Eveillard, D. (2024). Community metabolic modeling of host-microbiota interactions through multi-objective optimization. *Iscience, 27*(6).
- Li, W., Wang, L., Qiang, X., Song, Y., Gu, W., Ma, Z., & Wang, G. (2024). Design, construction and application of algae-bacteria synergistic system for treating wastewater. *Journal of Environmental Management, 366*, 121720.
- López-Palau, N. E., & Olais-Govea, J. M. (2020). Mathematical model of blood glucose dynamics by emulating the pathophysiology of glucose metabolism in type 2 diabetes mellitus. *Scientific reports, 10*(1), 12697.
- Lu, X. X., Zhu, Y. Y., Cheng, F., Wu, T., & Maiwulanjiang, M. (2024). Ajwain essential oil inhibits aflatoxin B1 and ergosterol production in *Aspergillus flavus*: Mechanisms and developmental stages-specific effects. *Food Bioscience, 62*, 105561.
- Luo, Y., Chen, Y., Gu, Z., Ni, R., Feng, P., Hu, Z., Song, L., Shen, X., Gu, C., & Li, J. (2023). Engineered muscle from micro-

- channeled PEG scaffold with magnetic Fe₃O₄ fixation towards accelerating esophageal muscle repair. *Materials Today Bio*, 23, 100853.
- March, S., Nerurkar, N., Jain, A., Andrus, L., Kim, D., Whittaker, C. A., Tan, E. K., Thiberge, S., Fleming, H. E., & Mancio-Silva, L. (2024). Autonomous circadian rhythms in the human hepatocyte regulate hepatic drug metabolism and inflammatory responses. *Science Advances*, 10(17), eadm9281.
- Pasricha, S.-R., Tye-Din, J., Muckenthaler, M. U., & Swinkels, D. W. (2021). Iron deficiency. *The Lancet*, 397(10270), 233-248.
- Shittu, S.-T. T., Lasisi, T. J., Shittu, S. A.-S., Adeyemi, A., Adeoye, T. J., & Alada, A. A. (2021). Ocimum gratissimum enhances insulin sensitivity in male Wistar rats with dexamethasone-induced insulin resistance. *Journal of Diabetes & Metabolic Disorders*, 20(2), 1257-1267.
- Taghilou, S., Peyda, M., & Mehrasbi, M. (2023). A review on the significance of Azolla for water and wastewater treatment. *Desalination and Water Treatment*, 293, 138-149.
- Ugolkov, Y., Nikitich, A., Leon, C., Helmlinger, G., Peskov, K., Sokolov, V., & Volkova, A. (2024). Mathematical modeling in autoimmune diseases: from theory to clinical application. *Frontiers in immunology*, 15, 1371620.
- Ulusoy-Gezer, H. G., & Rakicioğlu, N. (2024). The future of obesity management through precision nutrition: putting the individual at the center. *Current Nutrition Reports*, 13(3), 455-477.
- Wu, J., Zhang, B., Liu, X., Peng, L., Liu, J., Hu, Y., Ji, X., Lv, H., & Wang, S. (2023). Current gut-on-a-chip platforms for clarifying the interactions between diet, gut microbiota, and host health. *Trends in Food Science & Technology*, 134, 1-12.
- Zapata, F., Pastor-Ruiz, V., Ortega-Ojeda, F., Montalvo, G., Ruiz-Zolle, A. V., & Garcia-Ruiz, C. (2021). Human ultra-weak photon emission as non-invasive spectroscopic tool for diagnosis of internal states—A review. *Journal of Photochemistry and Photobiology B: Biology*, 216, 112141.
- Zhu, B., & Liu, S. (2023). Preservation of ~ 12-h ultradian rhythms of gene expression of mRNA and protein metabolism in the absence of canonical circadian clock. *Frontiers in Physiology*, 14, 1195001.