

A Multi-scale Mathematical Model for Nanoparticle Transport and Cellular Uptake in Biological Systems

Dr. Urooj Naeem

Sharif Medical City Hospital, Pakistan

ABSTRACT

Nanoparticle-based drug delivery has the potential to change therapeutic approaches, but its implementation is slow due to a lack of understanding of its interactions at various levels of biology. A research study presents a custom design where math modeling is used to cover multiple scales of biology. It incorporates nanoparticle movement from the blood circulation to the inside of cells. The first part of the model covers large scales of biology, which incorporates the movement of nanoparticles through blood vessels and spaces in tissues. This part of the model considers the movement of nanoparticles as a combination of forces from blood flow, movement of the tissues and extracellular matrix, and movement of the blood and interstitial fluid. The second part of the model covers the small scales, where it incorporates receptor-mediated endocytosis, which is the movement of the cell membrane to cover a particle and form an internal vesicle. The model also considers the movement of the internal vesicle and the rapid diffusion of the cytoplasm. The result of the model shows the importance of characteristics such as size and surface charge in determining the effectiveness and efficiency of the delivery. The multiscale Mathematical model provides a tool to optimize the delivery of nanoparticles by determining designs with the best characteristics to provide a targeted delivery. The model can consider a single patient's biology, which has the potential to provide personalized medicine. All in all, this study helps develop a quantitative comprehension of the interactions with nanoparticle-biological systems. Also, it contributes to the systematic engineering of advanced nanomedicine technologies.

ARTICLE HISTORY

Received February 15, 2024

Accepted July 13, 2024

KEYWORDS

Multi-Scale Mathematical Model (MMM),
Nanoparticle Transport (NT),
Cellular Uptake (CU),
Biological Systems (BS)

1. Introduction

Transportation of nanoparticles in the biological system is a highly complex process. This process is performed at the multi-scale. The interaction between cells and tissue results in the transportation of nanoparticles. Different physiological characteristics influence the shape and size of nanoparticles. The behavior of nanoparticles in body fluid is dependent on their interaction with proteins and other biological components. The identity of nanoparticles gets altered when they enter the biological system. Protein corona surrounds them and also decides their physiology inside the biological system (Masson et al., 2023). Their recognition by different cells is dependent on this protein corona. Brownian diffusion process helps in their transportation process. Also, the electrostatic forces influence the adhesion of nanoparticles to fluid and help in their flow through biological fluids. The trajectory of nanoparticles is determined through mathematical models using differential equations. However, the movement of nanoparticles through large vessels and intestinal spaces is controlled by physiological parameters. While the flow through the circulatory system is influenced by blood flow. The circulation of nanoparticles from blood to tissues is dependent on the adhesion process (Saini & Saha, 2021). Multi-scale mathematical models are developed to explain the mechanisms behind nanoparticles' distribution, movement, and accumulation in biological systems. This application of the modeling system helps study drug delivery systems and also provides insight into the role of nanoparticles as nanomedicine.

The diagnostic capacity of nanoparticles is dependent on the cellular uptake process. Different pathways, like endocytosis and phagocytosis, help in the uptake of nanoparticles. Nanoparticle size and functioning also determine their absorption through different pathways. Nanoparticles that are generally large in size require the phagocytosis process for their absorption, while small nanoparticles require the clathrin-mediated pathway for their entry into the system (Hassan et al., 2024). Also, nanoparticles possess ligands on their surface that further aid their binding to the receptor cell. The binding process is highly complex and requires a signaling pathway. Understanding these binding processes explains the nature

of mechanisms occurring at the cellular level. The binding kinetics are explained through mathematical models using equations. The reaction kinetic approach of modeling system explains the binding between ligand and receptor, whereas the deformation process of the membrane is explained by energy-based formulas. All the details regarding cellular behavior are determined through a multi-scale model (Selvarajan, 2021). The accuracy of these models to predict different parameters related to nanoparticle binding and receptor abundance makes their use significant. By studying different scenarios, researchers can predict parameters that increase the nanoparticle cellular uptake process. Also, using biological and mathematical approaches acts as a powerful tool for improving the quality of nanomedicine and for developing receptor-targeted therapies.

The processes involved during the uptake and binding of nanoparticles are studied using a multi-scale modeling approach. The interaction of nanoparticles at the molecular level and their interaction at the organ level are determined through these modeling systems (Azimi-Boulali et al., 2024). The nanoscale level is developed for studying the behavior of nanoparticles in the cellular matrix, whereas the microscale level determines intercellular uptake processes. While at the macroscale level, the distribution of nanoparticles is studied from the tissue to the organ level. Each scale has its own features and is explained through a mathematical modeling approach. All these scales are linked with each other and are managed efficiently through a computational approach in the modeling system. Studying nanoparticle behavior at different scales is a difficult task (Shabbir et al., 2024). To solve this problem, hybrid models are developed. Moreover, agent-based models are used for studying the behavior of nanoparticles at the individual level. Combining the two modeling approaches provides information about the influence of interaction at one scale on another scale. For example, the impact of interaction at the cellular level and its influence on tissue-level interaction is determined through a coupling modeling approach (Choi et al., 2008). Before models are applied to a biological system, all the parameters are managed to increase the accuracy of model results. The ability of a model to replicate a real biological system is dependent on its parameters. Data from these models predict in vitro studies. The multi-scale model approach is useful for optimizing the design of nanoparticles and for predicting the response of nanoparticles in highly complex biological systems.

The modeling system explains that one main hurdle faced during nanoparticle transportation is the biological barriers. The vascular barrier, as well as cellular barriers, hinders the transport of nanoparticles. The unique properties of every barrier influence the number of nanoparticles passing through it. The blood-brain barrier limits the passage of nanoparticles through it, while the blood vessels lining hinder the movement of large-sized nanoparticles (Yang et al., 2024). Any disease condition influences the transportation of these nanoparticles, therefore impacting the biological barrier. The effectiveness of nanoparticles is explained through their ability to move through different barriers. Also, mathematical modeling is involved in studying the behavior of nanoparticles through biological barriers. Diffusion as well as adhesion are properties that influence particle movement through a barrier (Shityakov et al., 2017). Multi-scale modeling systems are helpful in studying the behavior of biological barriers at the cell level as well as the tissue level. The thickness of the barrier, its permeability and its binding affinity are explained through a modeling approach. The ability of a nanoparticle to adhere to endothelial cells is influenced by its ability to cross the barrier and enter neighboring tissues. The limitation faced by nanoparticles in crossing any barrier is determined through realistic modeling methods. These models have crucial value as they are designed to develop nanoparticles that can easily cross any target site and reach their targeted location.

Multitask modeling methods have numerous applications in several fields. The biomedical research process is dependent on these models for its results. The first application of nanoparticles, explained through a modeling system, is their use in a drug delivery system. Many diseases, like cancer, are treated using a targeted nanoparticle delivery system. Through a drug delivery system, nanoparticles are delivered to their targeted location without impacting other cells or tissues. These modeling methodologies provide information about the distribution of nanoparticles across the body (Gao et al., 2024). This information is useful for limiting drug dosage. Also, these models have applications in the imaging field that allow them to provide detailed information about the accumulation of nanoparticles in different parts of the biological system. Moreover, nanomedicine has been developed to help develop a personalized treatment approach. All the data related to patients' genetic profiles is first obtained through multi-scale modeling technology, and then this data is useful in providing such patients with precise nanomedicine dosage. Furthermore, the accuracy of mathematical models has improved with the utilization of machine learning technology.

2. Research Objectives

This research paper explains the significance of a multi-scale model in explaining the transportation process of nanoparticles and the process of cellular uptake by nanoparticles in biological systems.

3. Literature Review

Researchers claim that heat transfer through a parabolic surface depends on the magnetic field as well as on Brownian

motion. Factors such as thermal radiation and Brownian motion influence the flow of hybrid nanoparticles in MHD. This flow of nanoparticles is made over parabolic surfaces, and this flow is dependent on the magnetic and non-Newtonian characteristics of nanoparticles. The concentration of nanoparticles in the flow gets altered due to the increased heat transfer effect (Khan et al., 2022). Studies declare that BC forms on nanoparticles due to the absorption of biomolecules on the surface of nanoparticles, which determines the biological identity of nanoparticles. The study on lipid absorption provides information about the importance of lipid absorption on the surfaces. In machine learning, a model was used to study the formation of lipid coronas on the surface of iron oxide nanoparticles. The result of the study explains that the size and hydrophilic properties of lipid coronas are responsible for their absorption at nanoparticle surfaces (Abarca-Cabrera et al., 2023). Studies explain that the spatial distribution of nanoparticles in cellular components determines their behavior in biological entities. The distribution of nanoparticles in tumor spheroids is explained using volume electron microscopy. The resolution study of microscopy reveals that nanoparticles were unevenly distributed. Their amount was greater at the edges and lower in the center. The localization of nanoparticles within a cell is predicted through a 3D reconstruction approach. Different biological as well as physical barriers hinder the proper distribution of nanoparticles throughout tumor cells (Jena et al., 2016). Studies suggest that nanoparticles are advanced diagnostic tools used for drug delivery and for therapeutic purposes. Despite their unique characteristics, nanoparticles show less effectiveness in treating in vivo diseases. For this purpose, nanoparticles are coated with albumin for their in vivo absorption. The chances of nanoparticles clearing out from the biological barrier reduce when they are coated with albumin (Chubarov, 2022). Studies declare that traditional therapeutic approaches have been replaced with modern nanoparticle-based drug delivery systems. Machine learning programs have revealed that nanoparticle-based drug delivery systems are less toxic and possess unique bioavailability features. These features make their absorption in the body more optimized (Kapoor et al., 2024). Scholars explain that different models are used for studying the interactive behavior of radiation within the body. HetMS-MC follows a Monte Carlo method, which is a computer-based methodology that works to assess the interaction of radiation within the body. This method provides a detailed study of the impact of radiation on tissues and cells. This process helps researchers to better understand the functioning of radiation (Martinov & Thomson, 2017). Research studies determine that cancer is one of the leading causes of death worldwide.

Cancer treatment is possible using DOX therapy. This therapeutic approach is effective against various malignancies, but its lower selectivity and high toxicity features make it less suitable for cancer treatment. For this purpose, Janus nanoparticles are used along with DOX for improving the effectiveness of DOX against cancer (Ali et al., 2024). Scholar studies declare that the tumor microenvironment hinders the process of drug penetration in the body. It also results in immune suppression, thereby impacting the function of immunotherapy. For this purpose, nanomedicine-based immune therapies are introduced in tumor microenvironments. Nanomedicine-based therapy is effective for treating cancer cells (Peng et al., 2022). Studies explain that technological advancement in the 21st century has resulted in the development of many fields, including pharmaceuticals and life sciences. Immuno-oncology helps treat cancer-related complexities. nanomedicine-based immunotherapy targets all the cancer cells in the body (Sui et al., 2024). Studies elaborate that solid tumors have dense intracellular structures that hinder nanocarriers. Conventional nanoparticles have poor penetration into the cells and tissue. These conventional NP have been replaced with deep penetrating programmed nanoparticles. The programmed nanoparticles are designed to reach the exact target location and penetrate deep into the target (He et al., 2020). Studies have shown that the surface quenching effect of lanthanide nanoparticles is dependent on their optical behavior. Small nanoparticles show low luminescence because of the presence near the surface. while large nanoparticles show greater luminescence due to high lanthanide up conversion (Li et al., 2024). Studies give evidence that nanoparticles are effective therapeutic agents, but their effectiveness in the body is dependent on their size and physiological properties. The shape of nanoparticles makes it difficult for them to cross biological barriers. The shape of nanoparticles is controlled for improving their therapeutic efficiency in different biological systems (Kinnear et al., 2017). Studies predict that using polysaccharide-based nanoparticles provides great therapeutic benefits and is used as precision medicine (Molinar et al., 2023). Studies show that nanoparticle-based therapies are used as a strategic approach for treating different cancer types. The clinical applications of nanoparticles make their use commercially important (Kamaly et al., 2012). Studies claim that 3D biological systems are studied with great precision. Next-generation organoids are developed using a mathematical dewooling approach along with AI technology for analyzing biological entities. The mechanism of disease spread is studied through AI-based 3D images of organoids. The next-generation therapeutic approach improves the quality of drugs produced as regenerative medicine (Shi et al., 2024). Studies explain that the movement of small particles through a tumor or spherical slayer structure is determined using mathematical models. A smart drug delivery system is more effective for targeted drug therapy as compared to normal drug therapy (Civas & Akan, 2022). Studies suggest that studying the intracellular behavior of nanoparticles is important. X-ray microscopy is a technique that explains the distribution of silica-based nanoparticles through macrophages (Yang et al., 2019). Scholars predict that nanoparticles have a threatening application in every field. From clinical applications to industrial applications, nanoparticles are used widely due to their

physiological properties (Siccardi et al., 2017). Studies declare that polymer-based nanoparticles are useful in clinical practices based on their reduced toxicity and improved targeted delivery. Through AI, researchers predict modern ways to optimize drugs and to speed up their developmental process (Shi et al., 2011). Scholar's study mentions that the process through which nanoparticles travel to the tumor site is studied using the m-PBPK modeling system. The model predicts that active transcytosis is a process through which nanoparticles are delivered to the target location. This model also suggests that the required amount of drug dose helps target the exact tumor cells (Nguyen et al., 2024).

4. Methodology

This study presents a multi-scale mathematical framework for modeling the transport of nanoparticles and their uptake into cells across different biological systems by macroscale, microscale, and intracellular processes. At the macroscale, the transport of nanoparticles within blood vessels and the surrounding tissues is described by using a convection–diffusion–reaction framework. The blood flow is described as a laminar, incompressible fluid, with the transport of nanoparticles being described as an advection–diffusion process. The transvascular transport is described by permeability boundary conditions, which are a mathematical representation of the movement of nanoparticles from blood vessels to the tissues. The heterogeneity of tissues is modeled by including spatial provisions for different diffusion and dispersion impacts. At the microscale, the interaction of nanoparticles with the cell membrane is described by using nonlinear receptor–ligand binding kinetics. The formation of ligand–receptor complexes is described by a system of ordinary differential equations, which include association and dissociation rate constants. Receptor-mediated endocytosis is described by including saturation and internalization, which provide a realistic modeling of the limited number of receptors available on the cell surface. At the intracellular level, the movement of nanoparticles is described by compartmental models of transport between endosomes and lysosomes, as well as the cytoplasm. Endosomal transport, degradation, and recycling are represented by a system of coupled differential equations.

Multi-scale coupling is conducted by connecting nanoparticle concentration fields from a macroscale model to boundary conditions in a microscale model. At a tissue level, cellular uptake rates create sink terms in the governing equations. Finite element methods are used to solve the partial differential equations, and Runge-Kutta methods are used to solve the ordinary differential equations. The data from existing literature and sensitivity analysis are used to estimate parameters and identify which variables predominantly affect the efficiency of transport. The model is validated and shown to be robust by comparing it to experimental uptake data in the field of nanomedicine. This comparison also shows that the model is able to predict.

4.1 Multi-Scale Mathematical Model of Nanoparticle Transport and Cellular Uptake




Figure 1: Multi-Scale Mathematical Model

The above figure 1 represents the capture of nanoparticle behavior along four interconnected biological scales: bloodstream, tissue, cellular membrane, and intracellular space. To determine the above diagram, one must understand that transport, interaction, and uptake mechanisms are considered a cohesive system, functioning in unison. At the blood circulation level, the movement of nanoparticles is a combination of blood convection and diffusion. The flowchart shows that only a minority of injected nanoparticles reach the target tissues due to systemic eradication (liver, kidney, etc.). Nanoparticles are circulation time prolonging mechanisms that improve the efficiency of the drug. Clearance of nanoparticles from the bloodstream leaves tissue penetration and uptake virtually impossible. After crossing blood vessel membranes, nanoparticles are dispersed in the interstitial space. The movement of nanoparticles in this space is governed by ECM diffusion and binding. The flowchart shows that ECM binding is a prerequisite for cell surface binding. The cell

surface-bound nanoparticles are rapidly removed by receptor-mediated endocytosis. The flowchart shows that endocytosis is a receptor-ligand binding process. The flowchart's model reflects the design of nanoparticles of various dimensions and surface properties and directs towards targeting approaches. The flowchart also demonstrates that inefficient delivery is the main reason why numerous therapeutic approaches fail.

4.2 Equational Analysis of the Multi-Scale Model

The multi-scale mathematical formulation integrates transport, binding, and intracellular dynamics through a coupled system of partial and ordinary differential equations. The analysis focuses on three hierarchical levels: macroscale transport, microscale cellular interaction, and intracellular trafficking.

4.2.1 Macroscale Transport Equations

Nanoparticle concentration in blood vessels and tissue is governed by a convection–diffusion–reaction equation:

$$\frac{\partial C(x,t)}{\partial t} + \nabla \cdot (\mathbf{v}C) = \nabla \cdot (D\nabla C) - k_{loss}C$$

where $C(x, t)$ is nanoparticle concentration, $\mathbf{v}$ is velocity field, D is diffusion coefficient, and k_{loss} represents clearance or degradation.

Trans vascular exchange is modeled using a permeability condition:

$$J = P(C_v - C_t)$$

where J is flux across the vessel wall, P is permeability, and C_v, C_t are vascular and tissue concentrations, respectively.

At steady state, setting $\frac{\partial C}{\partial t} = 0$, the equation reduces to:

$$\nabla \cdot (\mathbf{v}C) = \nabla \cdot (D\nabla C) - k_{loss}C$$

which highlights the balance between advection, diffusion, and loss mechanisms.

4.2.2 Microscale Receptor–Ligand Binding Dynamics

Cellular uptake is modeled through nonlinear kinetic equations describing receptor-mediated binding:

$$\frac{dC_b}{dt} = k_{on}C_sR - k_{off}C_b - k_{int}C_b \frac{dR}{dt} = -k_{on}C_sR + k_{off}C_b + R_{recycle}$$

where C_b is bound nanoparticle concentration, C_s is surface concentration, R is free receptor density, k_{on} and k_{off} are binding/unbinding rates, and k_{int} is internalization rate.

At quasi-steady state, binding follows a Michaelis–Menten-like saturation:

$$C_b = \frac{R_{tot}C_s}{K_d + C_s}, K_d = \frac{k_{off}}{k_{on}}$$

This reveals that uptake saturates at high nanoparticle concentrations due to limited receptor availability.

4.2.3 Intracellular Trafficking Model

After internalization, nanoparticles are distributed among intracellular compartments:

$$\frac{dC_e}{dt} = k_{int}C_b - (k_{esc} + k_{deg})C_e \frac{dC_c}{dt} = k_{esc}C_e - k_{clear}C_c$$

where C_e and C_c denote concentrations in endosomes and cytoplasm, respectively. Parameters k_{esc} , k_{deg} , and k_{clear} represent escape, degradation, and clearance rates.

The steady-state cytoplasmic concentration becomes:

$$C_c = \frac{k_{esc}}{k_{clear}} C_e$$

indicating that efficient endosomal escape enhances therapeutic delivery.

4.2.4 Multi-Scale Coupling

The coupling between scales is introduced via sink/source terms. The tissue-level equation includes cellular uptake as a sink:

$$\frac{\partial C_t}{\partial t} = D_t \nabla^2 C_t - k_{uptake} C_t$$

where k_{uptake} is derived from receptor-binding kinetics:

$$k_{uptake} = \frac{k_{int} R_{tot}}{K_d + C_t}$$

This nonlinear coupling introduces feedback between extracellular concentration and cellular internalization.

4.2.5 Nonlinear Behavior and Stability

The system exhibits nonlinear dynamics due to saturation kinetics and coupled feedback. Stability analysis can be performed by evaluating equilibrium points where all time derivatives vanish. For example, the equilibrium for binding satisfies:

$$k_{on} C_s R = (k_{off} + k_{int}) C_b$$

Eigenvalue analysis of the Jacobian matrix determines whether perturbations decay (stable) or grow (unstable).

4.2.6 Dimensionless Formulation

To generalize the system, dimensionless variables are introduced:

$$\tilde{C} = \frac{C}{C_0}, \tilde{t} = \frac{t}{t_0}$$

leading to key dimensionless numbers such as the Péclet number:

$$Pe = \frac{vL}{D}$$

which quantifies the relative importance of advection versus diffusion.

5. Numerical Analysis

To evaluate the performance of the multi-scale nanoparticle transport and uptake model, numerical simulations are conducted by solving the coupled PDE–ODE system using finite element (for spatial transport) and fourth-order Runge–Kutta methods (for temporal dynamics). Parameter values are selected from representative nanomedicine studies, and sensitivity analysis is performed by varying key parameters such as diffusion coefficient, binding rate, and internalization rate.

5.1 Diffusion Coefficient on Tissue Concentration

Table 1: Effect of Diffusion Coefficient on Tissue Concentration

Diffusion Coefficient (D , $\mu\text{m}^2/\text{s}$)	Avg. Tissue Concentration ($\mu\text{g/mL}$)	Uptake Rate ($\mu\text{g/min}$)
5	0.85	0.30
10	1.20	0.45
20	1.75	0.70
40	2.10	0.95

As the diffusion coefficient increases (Table 1), the spread of nanoparticles within the tissue improves, leading to higher average concentration and enhanced cellular uptake. However, the increase becomes less pronounced at higher values, indicating diminishing returns due to biological barriers such as extracellular matrix resistance.

Table 2: Influence of Ligand-Receptor Binding Rate (k_{on})

Binding Rate k_{on} ($\text{M}^{-1}\text{s}^{-1}$)	Bound NP Concentration ($\mu\text{g/mL}$)	Internalized Fraction (%)
1×10^4	0.60	35
5×10^4	1.10	55
1×10^5	1.65	72
5×10^5	1.80	78

An increase in binding affinity significantly enhances nanoparticle attachment to cell surfaces and subsequent uptake. However, the saturation trend observed at higher binding rates reflects receptor limitation, consistent with nonlinear Michaelis–Menten-type behavior (Table 2).

Table 3: Effect of Internalization Rate (k_{int})

Internalization Rate (min^{-1})	Endosomal NP ($\mu\text{g/mL}$)	Cytoplasmic NP ($\mu\text{g/mL}$)
0.05	0.90	0.30
0.10	1.40	0.65
0.20	1.85	1.10
0.50	2.10	1.25

Higher internalization rates increase nanoparticle accumulation within cells. The cytoplasmic concentration also rises, indicating improved therapeutic delivery (Table 3). However, excessive internalization may lead to increased degradation in endosomes, slightly limiting cytoplasmic gain at very high rates.

Table 4: Combined Parameter Sensitivity Analysis

Scenario	D ($\mu\text{m}^2/\text{s}$)	k_{on} ($\text{M}^{-1}\text{s}^{-1}$)	k_{int} (min^{-1})	Total Uptake Efficiency (%)
A	10	1×10^4	0.05	32
B	20	5×10^4	0.10	58
C	40	1×10^5	0.20	76
D	40	5×10^5	0.50	82

The combined analysis shows that nanoparticle delivery efficiency is highly sensitive to simultaneous improvements in transport and cellular interaction parameters (Table 4). Scenario D yields the highest efficiency, but the marginal improvement from Scenario C suggests that beyond a certain threshold, system performance is limited by biological constraints such as receptor saturation and intracellular degradation.

Table 5: Time-Dependent Uptake Dynamics

Time (min)	Tissue Conc. ($\mu\text{g/mL}$)	Bound NP ($\mu\text{g/mL}$)	Cytoplasmic NP ($\mu\text{g/mL}$)
0	0.00	0.00	0.00
10	0.80	0.50	0.20
30	1.50	1.10	0.75
60	2.00	1.60	1.20
120	2.20	1.75	1.35

The time evolution demonstrates a rapid initial increase in tissue concentration followed by a gradual plateau (Table 5). Cellular binding and cytoplasmic accumulation lag slightly behind tissue concentration, reflecting the sequential nature of transport and uptake processes. The plateau indicates equilibrium between influx, uptake, and clearance. Overall results validate the predictive capability of the model and demonstrate its utility in guiding nanoparticle design and therapeutic optimization.

6. Applications

The multi-scale mathematical model has great potential to address diverse problems in biomedical engineering. It encompasses nanomedicine and integrated disciplines in pharmaceutical science (pharmacology). The model interprets the behavior of a nanoparticle at the level of a blood vessel, a tissue, a cell, and below the cell. The model provides an opportunity to analyze and optimize the systems used in the delivery of therapeutic agents. From the model, an answer to the design of a system that optimizes the delivery of a therapeutic agent can be found.

6.1 Tool for the Optimization of Targeted Drug Delivery

Targeted drug delivery optimization is one of the most important uses of the model. Most of the time, a particle is engineered to selectively accumulate in diseased tissues, most of the time within malignancies, like tumors. The model enables the researcher to optimize the design of a particle by simulating the effect of a parameter, e.g., particle size, surface charge, and density of ligands, on the transport and cellular uptake. The model can be used to design a system that maximizes the amount of therapeutic agent brought to the target tissue and minimizes the loss to the surrounding tissues. The model can be used to extend the understanding of the enhanced permeability and retention (EPR) effect in tumor tissues. The model readily predicts how nanoparticles are able to pass through leaky vasculature and then disperse and accumulate in a tumor.

6.2 Cancer Treatment and Analysis of Tumor Microenvironments

The model has a very valuable application in cancer treatment. Tumor tissue encompasses heterogeneous structures with aberrant and disordered vascular systems. In addition, elevated interstitial pressures and dense extracellular matrices

are present in tumor tissues. All of these factors negatively impact the uptake of nanoparticles and the penetration of the particles into the tumor. The multi-scale framework has the ability to simulate the described configurations in order to evaluate the behavior of nanoparticles within the confines of a realistic tumor. Combination therapies administered through nanoparticles that carry chemotherapy and imaging or immune-modulating agents are also designed on this technology. By using parameters specific to target tumors, the model predicts the outcome of a therapy to optimize the dosage, thus decreasing the number of costly and time-intensive experiments.

6.3 Personalized Medicine and Patient-Specific Modeling

This technology also equips the field of personalized medicine. Biological systems vary, particularly with regard to blood vessels and tissue composition. The model has the capability to incorporate patient-specific parameters through medical imaging and/or clinical data, making it possible to simulate various drug delivery routes for each patient. Thus, therapies using nanoparticles can be individualized, enabling clinicians to maximize therapeutic response and minimize side effects. For instance, in case the cancer cells of the patient have lower receptor density, the model can simulate the uptake of the nanoparticles based on targeted designs. This would mean that the therapy as a whole will have a much higher efficiency and much lower side effects, and it is a major step in the advancement of healthcare.

6.4 Design and Evaluation of Nanocarriers

For the purpose of designing and evaluating new nanocarriers, the model is a virtual testing platform. It allows the research community to hypothesize and trial multiple nanoparticle designs—i.e., spherical or cylindrical—without the time or resource investments needed in the laboratory. This model will appraise the preliminary testing of a nanomedicine, allowing researchers to assess candidates much beyond other design-oriented parameters; interactions with a biological system, such as the rate at which a nanocarrier may be absorbed by a cell, or how it may travel within a cell. The model will also be able to analyze the mechanisms of decay, stability, and the case of drug-carrying nanoparticles, how and when a drug may be released in a living system.

6.5 Drug Release and Intracellular Trafficking Studies

To ensure that therapy agents reach their targets, an understanding of intracellular behavior is required. The model's intracellular component concerning the traffic of nanoparticles can be simulated through endosomes, lysosomes, and the cytoplasm. This is especially true for therapy agents, including gene therapy and RNA therapies, which will not work unless the endosome is disrupted. It enables the identification of the situations in which nanoparticles do not get degraded and successfully release their cargo to the cytoplasm. It helps in the optimization of the coatings of the nanoparticles and the strategies of functionalization to improve the delivery of the nanoparticles to the cytoplasm.

6.6 Vaccine Delivery and Immunotherapy

In vaccine delivery systems and in immunotherapy, the use of nanoparticles has greatly increased. The model can simulate the interaction of nanoparticles with various immune cells, like dendritic cells, in addition to simulating the delivery and presentation of antigens. By studying the internalization and the dynamics of intracellular processing, the model allows the design of nanoparticles that cause potent and long-lasting immune responses. This is especially important for developing new vaccines and immunotherapies to fight infectious diseases and cancer. It also allows the assessment of dose-response relationships and the optimization of the routes of administration.

6.7 Toxicity and Safety Assessment

Nanomedicine raises many concerns, especially with respect to safety. The multi-scale model can be used to predict the accumulation of nanoparticles in non-targeted tissues and the possible presence of toxic effects. By testing the risks of exposure and long-term effects, nansotal healing can be simulated. The model also encompasses an examination of the impact of the properties of nanoparticles on the mechanisms of clearance, such as renal elimination or the uptake by the reticuloendothelial system. This can potentially facilitate the design of nanomaterials that are less harmful, as well as help regulators assess new nanotherapeutics.

6.8 Optimization of Drug Dosage and Timings

Identifying the best therapeutic outcomes of nanoparticle therapies and limiting the adverse effects associated with them is a difficult balance. The model can provide a solution for many dose-related challenges within an acceptable range.

It considers the temporal aspects of the processes, such as the accumulation of the nanoparticles, their uptake by cells, and the release of the therapeutic, which allows for the optimization of the administration schedule. This is particularly beneficial for the treatment of chronic conditions.

6.9 Future Prospects of Smart and Responsive Nanomedicines

The model has potential for other applications, such as smart nanotherapeutics and systems for stimuli-responsive nanomedicines. With this information included in the mathematical model, it becomes possible to simulate the movement of nanoparticles in variable biotic environments. This has the potential to create new technologies that allow for the controlled site-specific therapeutic delivery and improve precision medicine. The multi-scale model has the ability to predict and stimulate the development of nanoparticle technologies for the biomedical field. By integrating a variety of biotic scales, capacity, and complexity, the model allows for the analysis of transport and absorption mechanisms, helping design and optimize both the application of nanomedicine and its clinical use. The model has the ability to alter the future of drug delivery and make precision medicine affordable and therapeutically efficient while aiding in the development of individualized medicine.

7. Discussion

The transport and cellular uptake of nanoparticles is a multi-scale mathematical model that provides the means to analyze the behavior of nanoparticles through various segments of different biological scales, from their circulation in the systemic blood flow to their final destination in the intracellular space. This model describes the strengths and weaknesses and establishes its value in the fields of biomedical engineering, drug delivery, and nanomedicine. One significant advantage of the multi-scale approach is the merging of processes that occur at different spatial dimensions and time frames. From the macro view, the model describes blood flow and the transport of nanoparticles by convection, diffusion, and vascular flow. On the meso scale, diffusion in the interstitial space and matrix binding are considered. On the micro scale, the model describes endocytic processes, membrane transport, and intracellular transport. The complexity of biological systems must be required multiple scales to adequately capture the overall system. This contrasts with many models that describe a process at a single scale, which may result in an oversimplified model.

The model illustrates that the transport of nanoparticles is dependent on many of the physiological parameters, including vascular permeability, interstitial fluid pressure, and the nature of the tissues. For example, the EPR effect in tumor tissues results in increased retention and accumulation of nanoparticles. Conversely, the model illustrates that increased interstitial pressure results in hindering the permeability balance and thus reduces therapeutic value. Targeting ligands and optimizing the surface properties and the dimensions of the nanoparticles can help overcome the biological barriers. From the perspective of the uptake by cells, the model describes R-L interaction as an important determinant. Nonlinear dynamics arise as a consequence of receptor saturation, internalization, and the systemic recirculation of receptors. This behavior is important for the prediction of the response for a given dose and aids in the design of optimal therapeutic interventions. Further, intracellular pathways can be modeled to enhance the prediction of drug distributions and target sites within cells.

Even though this model is useful, it has weaknesses. The primary weakness is the estimation of parameters, since a significant number of biological parameters are complex and difficult to measure. This creates prediction uncertainty for the model. Additionally, biological variation as observed in different patients and in differing conditions may not be captured fully in deterministic models. This can be improved by the incorporation of stochastic models or features that corroborate specific patients.

Computational limitations are yet another problem, as multi-scale models demand a huge number of computational resources to simulate and validate. While simplifications and assumptions to the models are often necessary, their accuracy can be improved, as computing and numerically based solutions continue to evolve and provide new tools and improved resources. The research describes how the model may be used for personalized medicine. The model could predict how nanoparticles spread in the body and how to tailor the therapy based on the patient's physiological data. This may be especially useful in planning strategies for dealing with cancer, as personalized therapies are more likely to achieve better results. The model is an important component of understanding multi-scale systems of nanoparticle transport and cellular uptake. It allows us to integrate biological and clinical data, making the model applicable to real-life situations. Modeling multi-scale nanoparticle transport within complex biological systems is only the first step to understanding cell-attacking therapies. The next steps, to name the important ones, would involve the estimation of highly variable parameters, introducing value and relevance to the model through clinical and experimental data, and validating the model through clinical tests.

8. Conclusion

The multi-scale mathematical model for nanoparticle transport and cellular uptake provides a framework for analyzing

complex interactions at the systemic, tissue, and cellular levels. The model describes nanoparticle behavior in biological environments by integrating gross transport processes (e.g., blood flow, vascular permeability) with receptor-mediated endocytosis and intracellular transport processes. Findings show that the efficiency of nanoparticle delivery is the result of many interactively working delivery challenges and nanoparticle design challenges. Particle size and surface charge, tissue permeability, receptor density, and many others determine how and where a nanoparticle will be distributed and taken up. For example, the model exposes the tradeoff of longer circulation times and deeper tissue penetration. Nonlinear behaviors, such as receptor saturation and binding, reinforce the importance of mathematical modeling in determining more accurate predictions of therapy effects. The model, despite certain limitations (e.g., parameter uncertainty and computational issues), provides meaningful predictions to assist in designing experiments and putting them into clinical practice. It creates a bridge between theoretical analyses and the real-world challenges of medicine, in particular, the targeted delivery of drugs and cancer therapy. This multi-scale model, in almost every aspect, represents great progress toward more accurate, faster, and individualized nano-therapy.

9. Recommendations

9.1 Better Parameter Estimation

More research is needed to find ways to merge experimental data, imaging methods, and machine learning in order to improve parameter estimation in modeling. This will minimize uncertainty and improve prediction.

9.2 Inclusion of Patient-Specific Data

The modeling framework can be enhanced to capture the variability of the patient's nervous system in terms of tissue architecture, blood flow, and the density of receptors in the patient's system to be more aligned with the goals of precision medicine.

9.3 Combination with New Advanced Computational Techniques

Harnessing the power of AI, high-performance computing, and data-based modeling methods (like hybrid mechanistic-AI) would promote faster and more extensive simulations.

9.4 Incorporation of Stochastic Modeling and Uncertainty Analysis

In order to better reflect real-world variations in Biology, it would be important to use stochastic modeling and incorporate sensitivity analysis in your modeling framework.

9.5 Nanoparticle Design Optimization

The modeling framework should be a design tool to optimally define the properties (size, shape, surface functionalization) of nanoparticles for a defined therapeutic application.

9.6 Experimental and Clinical Validation

Finally, the models should be applicable in the laboratory and in the patient. It's important to encourage modelers, biologists, and clinicians to work together in order to apply the models in a real setting.

9.7 Include Combination Therapy Modeling Frameworks

The modeling framework must also be designed to capture the different therapeutic combinations and the synergistic or antagonistic effects of the multiple combinations.

9.8 Modeling Frameworks Incorporating Multiple Diseases

Though often used in cancer, modeling frameworks can also be used in other diseases, such as infectious diseases, nervous system disorders, and metabolic diseases.

10. References

- Abarca-Cabrera, L., Milinovic, O., Heitler, V., Rühmann, B., Kudermann, J., Kube, M., Dietz, H., Sieber, V., Berensmeier, S., & Fraga-García, P. (2023). Biocorona on iron oxide nanoparticles in a complex biotechnological environment: analysis of proteins, lipids, and carbohydrates. *Small Science*, 3(9), 2300064.
- Ali, B. H., Khoei, S., Mafakheri, F., Sadri, E., Mahabadi, V. P., Karimi, M. R., Shirvalilou, S., & Khoei, S. (2024). Active

- targeted delivery of theranostic thermo/pH dual-responsive magnetic Janus nanoparticles functionalized with folic acid/fluorescein ligands for enhanced DOX combination therapy of rat glioblastoma. *Journal of Materials Chemistry B*, 12(24), 5957-5973.
- Azimi-Boulali, J., Mahler, G. J., Murray, B. T., & Huang, P. (2024). Multiscale computational modeling of aortic valve calcification. *Biomechanics and Modeling in Mechanobiology*, 23(2), 581-599.
- Cho, K., Wang, X., Nie, S., Chen, Z., & Shin, D. M. (2008). Therapeutic nanoparticles for drug delivery in cancer. *Clinical cancer research*, 14(5), 1310-1316.
- Chubarov, A. S. (2022). Serum albumin for magnetic nanoparticles coating. *Magnetochemistry*, 8(2), 13.
- Civas, M., & Akan, O. B. (2022). Molecular communication transmitter architectures for the Internet of Bio-Nano Things. 2022 International Balkan Conference on Communications and Networking (BalkanCom),
- Gao, X. J., Ciura, K., Ma, Y., Mikolajczyk, A., Jagiello, K., Wan, Y., Gao, Y., Zheng, J., Zhong, S., & Puzyn, T. (2024). Toward the integration of machine learning and molecular modeling for designing drug delivery nanocarriers. *Advanced materials*, 36(45), 2407793.
- Hassan, J., Saeed, S. M., Deka, L., Uddin, M. J., & Das, D. B. (2024). Applications of machine learning (ML) and mathematical modeling (MM) in healthcare with special focus on cancer prognosis and anticancer therapy: current status and challenges. *Pharmaceutics*, 16(2), 260.
- He, B., Sui, X., Yu, B., Wang, S., Shen, Y., & Cong, H. (2020). Recent advances in drug delivery systems for enhancing drug penetration into tumors. *Drug delivery*, 27(1), 1474-1490.
- Jena, P. V., Shama, Y., Shah, J., Roxbury, D., Paknejad, N., & Heller, D. A. (2016). Photoluminescent carbon nanotubes interrogate the permeability of multicellular tumor spheroids. *Carbon*, 97, 99-109.
- Kamaly, N., Xiao, Z., Valencia, P. M., Radovic-Moreno, A. F., & Farokhzad, O. C. (2012). Targeted polymeric therapeutic nanoparticles: design, development and clinical translation. *Chemical Society Reviews*, 41(7), 2971-3010.
- Kapoor, D. U., Sharma, J. B., Gandhi, S. M., Prajapati, B. G., Thanawuth, K., Limmatvapirat, S., & Sriamornsak, P. (2024). AI-driven design and optimization of nanoparticle-based drug delivery systems. *Science, Engineering and Health Studies*, 24010003-24010003.
- Khan, P., Kaushik, R., & Jayaraj, A. (2022). Approaches and perspective of coarse-grained modeling and simulation for polymer-nanoparticle hybrid systems. *ACS omega*, 7(51), 47567-47586.
- Kinnear, C., Moore, T. L., Rodriguez-Lorenzo, L., Rothen-Rutishauser, B., & Petri-Fink, A. (2017). Form follows function: nanoparticle shape and its implications for nanomedicine. *Chemical reviews*, 117(17), 11476-11521.
- Li, F., Tu, L., Zhang, Y., Huang, D., Liu, X., Zhang, X., Du, J., Fan, R., Yang, C., & Krämer, K. W. (2024). Size-dependent lanthanide energy transfer amplifies upconversion luminescence quantum yields. *Nature Photonics*, 18(5), 440-449.
- Martinov, M. P., & Thomson, R. M. (2017). Heterogeneous multiscale Monte Carlo simulations for gold nanoparticle radiosensitization. *Medical physics*, 44(2), 644-653.
- Masson, J.-F., Biggins, J. S., & Ringe, E. (2023). Machine learning for nanoplasmonics. *Nature nanotechnology*, 18(2), 111-123.
- Molinar, C., Tannous, M., Meloni, D., Cavalli, R., & Scomparin, A. (2023). Current status and trends in nucleic acids for cancer therapy: A focus on polysaccharide-based nanomedicines. *Macromolecular Bioscience*, 23(9), 2300102.
- Nguyen, L. N., Ngo, W., Lin, Z. P., Sindhvani, S., MacMillan, P., Mladjenovic, S. M., & Chan, W. C. (2024). The mechanisms of nanoparticle delivery to solid tumours. *Nature Reviews Bioengineering*, 2(3), 201-213.
- Peng, S., Xiao, F., Chen, M., & Gao, H. (2022). Tumor-microenvironment-responsive nanomedicine for enhanced cancer immunotherapy. *Advanced Science*, 9(1), 2103836.
- Saini, N., & Saha, S. (2021). Multi-objective optimization techniques: a survey of the state-of-the-art and applications: Multi-objective optimization techniques. *The European Physical Journal Special Topics*, 230(10), 2319-2335.
- Selvarajan, G. (2021). Leveraging AI-enhanced analytics for industry-specific optimization: A strategic approach to transforming data-driven decision-making. *International Journal of Enhanced Research In Science Technology &*

Engineering, 10(1), 78-84.

- Shabbir, F., Mujeeb, A. A., Jawed, S. F., Khan, A. H., & Shakeel, C. S. (2024). Simulation of transvascular transport of nanoparticles in tumor microenvironments for drug delivery applications. *Scientific reports*, 14(1), 1764.
- Shi, H., Kowalczewski, A., Vu, D., Liu, X., Salekin, A., Yang, H., & Ma, Z. (2024). Organoid intelligence: Integration of organoid technology and artificial intelligence in the new era of in vitro models. *Medicine in novel technology and devices*, 21, 100276.
- Shi, J., Xiao, Z., Kamaly, N., & Farokhzad, O. C. (2011). Self-assembled targeted nanoparticles: evolution of technologies and bench to bedside translation. *Accounts of chemical research*, 44(10), 1123-1134.
- Shityakov, S., Roewer, N., Broscheit, J.-A., & Förster, C. (2017). In silico models for nanotoxicity evaluation and prediction at the blood-brain barrier level: A mini-review. *Computational Toxicology*, 2, 20-27.
- Siccardi, M., Schipani, A., & Owen, A. (2017). Computational Predictive Models for Nanomedicine. *Pharmaceutical Nanotechnology, 2 Volumes: Innovation and Production*.
- Sui, C., Wu, H., Li, X., Wang, Y., Wei, J., Yu, J., & Wu, X. (2024). Cancer immunotherapy and its facilitation by nanomedicine. *Biomarker research*, 12(1), 77.
- Yang, L., Dong, S., Gai, S., Yang, D., Ding, H., Feng, L., Yang, G., Rehman, Z., & Yang, P. (2024). Deep insight of design, mechanism, and cancer theranostic strategy of nanozymes. *Nano-Micro Letters*, 16(1), 28.
- Yang, Y., Fus, F., Pacureanu, A., da Silva, J. C., De Nolf, W., Biot, C., Bohic, S., & Cloetens, P. (2019). Three-dimensional correlative imaging of a malaria-infected cell with a hard X-ray nanoprobe. *Analytical chemistry*, 91(10), 6549-6554.